

***Final Draft***  
**of the original manuscript:**

Zou, J.; Wang, W.; Kratz, K.; Xu, X.; Nie, Y.; Ma, N.; Lendlein, A.:  
**Evaluation of human mesenchymal stem cell senescence, differentiation and  
secretion behavior cultured on polycarbonate cell culture inserts.**  
In: Clinical Hemorheology and Microcirculation. Vol. 70 (2019) 4, 573 - 583.  
First published online by IOS: February 22, 2019

DOI: 10.3233/CH-189322  
<https://dx.doi.org/10.3233/CH-189322>

## **Evaluation of human mesenchymal stem cell senescence, differentiation and secretion behavior cultured on polycarbonate cell culture inserts**

Jie Zou <sup>1,2,#</sup>, Weiwei Wang <sup>1,#</sup>, Karl Kratz <sup>1,3</sup>, Xun Xu <sup>1,2</sup>, Yan Nie <sup>1</sup>, Nan Ma <sup>1,2,3,\*</sup> and Andreas Lendlein <sup>1,2,3,\*</sup>

<sup>1</sup> Institute of Biomaterial Science and Berlin-Brandenburg Center for Regenerative Therapies, Helmholtz-Zentrum Geesthacht, Kantstraße 55, 14513 Teltow, Germany

<sup>2</sup> Institute of Chemistry and Biochemistry, Freie Universität Berlin, Takustraße 3, 14195 Berlin, Germany

<sup>3</sup> Helmholtz Virtual Institute “Multifunctional Biomaterials for Medicine”, Kantstraße 55, 14513 Teltow, Germany

<sup>#</sup>These authors contributed equally to this work.

Corresponding authors: Nan Ma, E-Mail: nan.ma@hzg.de and Andreas Lendlein, E-Mail: andreas.lendlein@hzg.de

### **Abstract**

Polycarbonate (PC) substrate is well suited for culturing human mesenchymal stem cells (MSCs) with high proliferation rate, low cell apoptosis rate and negligible cytotoxic effects. However, little is known about the influence of PC on MSC activity including senescence, differentiation and secretion. In this study, the PC cell culture insert was applied for human MSC culture and was compared with polystyrene (PS) and standard tissue culture plate (TCP). The results showed that MSCs were able to adhere on PC surface, exhibiting a spindle-shaped morphology. The size and distribution of focal adhesions of MSCs were similar on PC and TCP. The senescence level of MSCs on PC was comparable to that on TCP, but was significantly lower than that on PS. MSCs on PC were capable of self-renewal and differentiation into multiple cell lineages, including osteogenic and adipogenic lineages. MSCs cultured on PC secreted a higher level inflammatory cytokines and pro-angiogenic factors including FGF2 and VEGF. Conclusively, PC represents a promising cell culture material for human MSCs.

**Key words:** Polycarbonate, human mesenchymal stem cells, differentiation, cytokine secretion, senescence

## **1. Introduction**

Mesenchymal stem cells (MSCs) attracted interest because of their promising therapeutic potential in regenerative medicine due to their self-renewal, multi-lineages differentiation potential and the secretion of a variety of functional cytokines [1-3]. MSCs can be obtained from different tissues, such as bone marrow, adipose tissue and umbilical cord tissue [4-6]. However, the quantity of MSCs that separated from those tissues can not satisfy the needs of clinic cell therapy (10–400 million MSCs per treatment) [7]. Therefore, the isolated MSCs need to be expanded *in vitro* before implantation. It has been shown that MSCs are very sensitive to the substrate properties, such as the extracellular matrix (ECM) stiffness [8], surface topography [9] and chemical property of the culture surface [10]. The functions and behavior of MSCs can be influenced during the period of cell expansion *in vitro* [11].

Polycarbonate (PC) has technological significance in life science applications, such as intravenous connection components or cell culture devices [12, 13]. Favorable properties of PC include the high light transmittance, impact toughness, dimensional stability, creep resistance and thermoplastic process ability [14]. In addition, PC can be sterilized by a variety of methods including ethylene oxide (EtO), irradiation and steam autoclaving. As cell culture devices, PC exhibited excellent cell compatibility, favoring cell adhesion and proliferation [15]. However, there are few reports about the influence of PC on the most important behaviors of MSCs such as senescence, differentiation and secretion.

Here we hypothesized that PC is a suitable substrate for cultivation of human bone marrow mesenchymal stem cells (hBMSCs) without further modification. In this study we compare a PC-based substrate with polystyrene-based tissue culture plate (TCP), whose surface is modified in order to support adherent cells, and a polystyrene (PS)-based substrate without surface modification. Those substrates serve as positive and negative reference materials. The cell adhesion, morphology, senescence, cytokine secretion and differentiation of hBMSCs on PC, PS and TCP were compared.

## **2. Materials and methods**

### *2.1. Materials for cell culture*

The polycarbonate (PC, trade name Makrolon® 2805, Bayer, Germany) and polystyrene (PS, Type 158K, BASF, Germany) cell culture inserts were fabricated via injection molding [16]. The surfaces of PS and PC inserts showed the similarity in both mechanical properties and surface profile, as reported previously [15]. The Young's modulus of PC and PS was  $13 \pm 2$  and  $16 \pm 4$  GPa respectively, as measured with AFM indentation. Profilometry analysis at an area of  $7 \times 7$  mm<sup>2</sup> indicated that the bottom roughness of the inserts is at a similar level (root arithmetic mean square roughness (Rq) =  $0.12 \pm 0.04$  µm for PS and  $0.34 \pm 0.13$  µm for PC). In addition, the PS and PC presented comparable surface wettability (water advancing contact angle  $85 \pm 8^\circ$  for PC and  $99 \pm 5^\circ$  for PS), as determined via water contact angle measurement using the captive bubble method [15]. Prior to cell culture, the PC and PS inserts were sterilized via gas sterilization (gas phase: 10% (v/v) ethylene oxide, 54 °C, 65% relative humidity, 1.7 bar, 3 hours of gas exposure time and 21 hours of aeration phase). In addition, the tissue culture plate (TCP, Corning, New York, USA) was used in this study as a positive control material.

## *2.2. Human bone marrow mesenchymal stem cells (hBMSCs)*

The hBMSCs were purchased from Merck Millipore (SCC034, Merck Millipore, Darmstadt, Germany). The cells were maintained in MesenPRO RS™ growth medium (ThermoFisher Scientific, Waltham, USA), in a humidified atmosphere containing 5% (v/v) CO<sub>2</sub>. The medium was changed in every 3 days. For hBMSC phenotype characterization, the cells were harvested and stained with a Human MSC Analysis Kit (Stemflow™, BD Biosciences, Heidelberg, Germany) according to the manufacturer's instruction. A flow cytometer (MACSQuant®, Miltenyi Biotec, Bergisch Gladbach, Germany) was used to analyze the cells and the data was processed using Flowjo software (Tree Star Inc., Ashland OR, USA).

## *2.3. Immunostaining*

Cells were fixed with 4% (w/v) paraformaldehyde (Sigma–Aldrich, St. Louis, MO, USA) and permeabilized with 0.1% (v/v) Triton X-100 (Sigma–Aldrich, St. Louis, MO, USA), then blocked with 3% (w/v) BSA buffer. Vinculin was stained with mouse anti-human vinculin monoclonal antibody (Merck Millipore, Darmstadt, Germany) and Alexa Fluor® 488 labeled IgG antibody (Life Technologies, Darmstadt, Germany). F-actin was stained with ActinRed™

555 (Life Technologies, Darmstadt, Germany). After washing, the cells were visualized with a confocal laser-scanning microscope (LSM 780 META, Carl Zeiss, Jena, Germany).

#### *2.4. Senescence assay*

The senescence-associated  $\beta$ -galactosidase activity of hBMSCs cultured on different materials was quantified using a cellular senescence assay kit (Cell Biolabs, Inc., San Diego, CA, USA) according to the manual introduction at different time points (day 1, 8, 15 and 22). The concentration of the total protein in the cell extraction was determined using a BCA protein assay kit (Thermo Fisher Scientific, Bonn, Germany). The cellular senescence level was expressed as the fluorescent intensity normalized with the amount of total protein.

#### *2.5. Secretion assay*

The cytokines in the conditioned medium and the cell number were quantified at day 4, 7, 10, 13, 16 and 19. The cytokines in the conditioned medium were quantified using a Bio-Plex® 200 system (BioRad, Munich, Germany) according to the manufacturer's instructions. The cell numbers at different time points were determined using a Cell Counting Kit-8 (CCK-8, Dojindo Molecular Technologies, Munich, Germany) as described before [15]. To evaluate the activity of cell secretion, the cytokine amount was normalized with the cell number.

#### *2.6. Differentiation assay*

For the differentiation study, the hBMSCs were cultured in osteogenic (StemPro Osteogenesis differentiation kit) and adipogenic (StemPro Adipogenesis differentiation kit) induction media, respectively. The induction media were changed in every 3 days. To evaluate the adipogenic differentiation, the cells were cultured in adipogenic induction medium for 14 days. Then, the cells were fixed with 4% (w/v) paraformaldehyde and permeabilized with 0.1% (v/v) Triton X-100 solution. After blocking with 3% (w/v) BSA, staining was performed using rabbit anti-human FABP-4 monoclonal antibody (Merck Millipore, Darmstadt, Germany) and Alexa Fluor® 488 labeled IgG antibodies (Life Technologies, Darmstadt, Germany). Cell nuclei were stained with SYTOX Green Nucleic Acid Stain (Life Technologies, Darmstadt, Germany). The stained samples were scanned with a confocal laser scanning microscope (LSM 780, Carl Zeiss,

Jena, Germany). Osteogenic differentiation was evaluated after culturing the cells in osteogenic induction medium for 21 days. The cells were fixed with 4% (w/v) paraformaldehyde, and Alizarin red S (Sigma–Aldrich, St. Louis, MO, USA) staining was performed to detect the mineral deposition.

## 2.7. Statistics

The number of replication for quantitative experiments was three as indicated in figure legend, and the data were expressed as arithmetic mean  $\pm$  standard deviation (SD). Statistical analysis was performed by one-way ANOVA with *post hoc* Tukey HSD test. A *p* value less than 0.05 was considered statistically significant.

## 3. Results

### 3.1. hBMSC phenotype

To analyze the phenotype of the cells, the cell morphology and surface markers were characterized. The cells presented the typical MSC's spindle-shaped morphology (Fig. 1A). Furthermore, the cell surface markers were analyzed with flow cytometry. The results indicated that the typical MSC markers (CD44, CD73, CD90 and CD105) were well preserved in the cells and the non-MSC markers (CD45, CD34, CD11b, CD19 and HLA-DR) were not detected (Fig. 1 B- F).




Fig. 1: Characterization of hBMSCs. A: The image of hBMSCs growing on TCP. B, C, D, and E: characterization of MSC phenotypic markers (CD44, CD73, CD90 and CD105). F: characterization of non-MSC markers (CD45, CD34, CD11b, CD19 and HLA-DR).

### 3.2. Immunostaining of focal adhesion complex

To study the cell-material interaction, the formation of focal adhesions (FAs) was examined by immunofluorescent staining of vinculin and F-actin (Fig. 2). As hBMSCs were difficult to attach and spread on PS surface, the focal adhesion (FA) formation of cells on PC and TCP surfaces were characterized. After 4 days of cultivation in growth medium, the cells spread out on the PC and TCP. The F-actin cytoskeleton organization and FA complex formation were similar in cells growing on PC and TCP.




Fig. 2: Representative laser scanning microscopic images of hBMSCs cultured on PC and TCP for 4 days. F-actin (red) and vinculin (green) were fluorescently stained. (scale bar = 20  $\mu\text{m}$ ).

### 3.3. Senescence of hBMSCs

The influence of culture materials on hBMSC senescence level was evaluated by measuring the activity of senescence-associated  $\beta$ -galactosidase (SA- $\beta$ -gal). The results indicated that in the early stage (day 1) the senescence level of hBMSCs on TCP surface was significantly higher than that on PS. With the elongation of cultivation, the senescence level increased on all of the tested surfaces. The senescence level of cells on PS was higher than that on PC and TCP at day 8 and day 22. There was no significant difference of the senescence level between the cells cultured on PC and TCP (Fig. 3).




Fig. 3: The senescence level of hBMSCs cultured on different materials at the indicated time points. (n = 3; \*p < 0.05).

#### 3.4. Cytokine secretion

To analyze the paracrine factors secreted by hBMSCs on different materials, the conditioned media of hBMSC cultures were collected and the factors were quantified (Fig. 4). The paracrine factors secretion level of cells on PS is higher than that on TCP and PC. In the early stage (day 4) the cells on PC showed a higher secretion level than the cells on TCP except for the MCAF secretion. Compared to TCP, the cells on PC secreted the higher level of inflammatory cytokines such as IL-6, IL-17, IFN- $\gamma$  and TNF- $\alpha$  as well as pro-angiogenic factors including FGF2 and VEGF.




Fig. 4: The heatmap showed the fold change (compared to TCP) of protein secretion over time by hBMSCs seeded on different materials.

### 3.5. Differentiation of hBMSCs

The osteogenic and adipogenic differentiation of hBMSCs on different materials were examined by Alizarin Red S (ARS) staining and FABP-4 immunostaining, respectively. The results indicated that hBMSCs did not differentiate in the non-induced medium on all of the tested surfaces (Fig. 5). When cultured in osteogenic induction medium, the cells could differentiate into osteoblasts (Fig. 5A). In addition, the immunofluorescence staining of FABP-4 indicated that the hBMSCs could be induced for adipogenic differentiation when adipogenic induction medium was applied (Fig. 5B).




Fig. 5: Differentiation of hBMSCs. A: The representative images of alizarin red S (ARS) staining for characterization of hBMSC osteogenic differentiation on different materials (bar = 100 μm). B: The representative laser scanning microscopic images of FABP-4 immunostaining for characterization of hBMSC adipogenic differentiation (red: FABP-4; blue: nuclei; bar = 100 μm).

#### 4. Discussion

The cell behavior can be regulated by substrate properties, such as the surface free energy, the surface polarity [17], the presence of functional groups [18], the stiffness of substrate [19] and the surface topography [20]. The cells actively sense and react to the substrates through the activation of cell surface integrin and then the focal adhesion (FA) complex formation [21-23]. The cell surface integrin activation can further activate the focal adhesion kinase (FAK), which is indispensable for the focal adhesion complex formation [24]. The FA complex consists of many signaling molecules, including the vinculin, talin, paxillin, Src and Cas [25-27]. These signaling molecules can be regulated by chemical and physical properties of the substrates, and

play a key role for remodeling the size, shape and distribution of FAs [28]. The FAs can be found at the peripheral and central areas of the cells. They are linked with the terminal of F-actin to regulate cells via transfer the signaling from the outside to inside [29]. Here, the formation and distribution of FAs were characterized by staining F-actin and vinculin. The images showed that the distribution and size of FAs on PC were comparable to that on TCP, indicating the excellent cell compatibility of PC for hBMSC adhesion. This observation was in line with our previous results [15].

MSCs are multipotent cells, which are capable of extensively replicating and maintaining their multi-potential during culture. However, the long-term cell cultivation *in vitro* may lead to cellular aging, alteration of genetic and epigenetic features as well as spontaneous cell transformation [30]. The primary cells do not grow infinitely, but undergo only a limited number of cell division, in a process called cellular senescence [31]. The MSCs cease growth *in vitro* at about 40 to 50 cell doublings [32]. The stem cell senescence could result in impaired regenerative capacity and reduced tissue function, which might consequently lead to organismal aging *in vivo* [33-36].

The senescent cells undergo irreversible growth arrest, but the metabolism is active. They normally present a large and flat morphology and display the characteristic changes in gene expression. Senescent cells typically exhibit an increase of SA- $\beta$ -gal activity [37, 38]. In this study, the result showed that with the elongation of cultivation the senescence level of hBMSCs increased on all of the tested materials, which could be explained by the regulation from the genetic factors. The senescence level of hBMSCs on PS was significantly higher than that on PC and TCP, which might be caused by the poor cell adhesion and spreading of cells on PS [15].

In addition to growth arresting, morphology change and accumulation of lipofuscin in cell plasma, senescence in stem cells could cause the damage of DNA and the alteration of functions [37]. In senescent MSCs, the potential of differentiation decreased with the disruption of the balance between osteogenic and adipogenic differentiation [39-41]. Moreover, long-term cultivation of MSCs could result in the complete loss of osteogenic potential and the decrease of adipogenic potential [42]. In this study, similar to hBMSCs on TCP, cells cultured on PC could be induced to differentiate towards osteogenic and adipogenic lineages by induction medium, and kept in undifferentiated status in growth medium. These results demonstrated that the hBMSCs were capable of maintaining their self-renewing and differentiation potential on PC.

MSCs can secrete a broad range of paracrine factors [1, 2, 43, 44]. These factors could exert the anti-inflammatory, anti-apoptotic, anti-fibrotic and pro-angiogenic functions [45]. The secretion of MSCs could be modulated by various cues from the microenvironment, such as the extracellular matrix (ECM) stiffness [8], surface topography[9] and the chemical property of substrate surface [10]. In our study, the hBMSCs on PS showed higher secretion activity compared to that on PC and TCP, which might be due to the poor cell adhesion and low cell number on PS. Notably, the higher secretion activity of inflammatory cytokines and pro-angiogenic factors including FGF2 and VEGF was observed in cells on PC, as compared to those on TCP. Since VEGF is one of the most important growth factors for promoting angiogenesis [46], this observation suggested that the pro-angiogenic capacity of MSCs might be improved by cultivation on PC.

In summary, our results demonstrated the high cell biocompatibility of PC for stem cell cultivation. The hBMSCs could attach on PC, forming FAs and cytoskeleton similar to those on TCP. Their self-renewal and differentiation potential could be maintained. Compared to TCP, hBMSCs on PC presented a comparable level of senescence and a higher secretion activity.

## **5. Conclusion**

In this study, the hBMSC senescence level, secretion behavior and differentiation potential on PC were evaluated comparing cell culture on TCP and PS. The PC showed high cell compatibility for hBMSC adhesion and spreading. Cells on PC presented the similar focal adhesion and cytoskeletal structure, as compared to TCP. The cellular senescence of hBMSCs on PC was at a comparable level to that on TCP. Cells on PC showed a higher secretion activity than on TCP, with the upregulated inflammatory cytokines and pro-angiogenic factors. The hBMSCs could maintain their stemness in growth medium on PC, and could undergo osteogenic and adipogenic differentiation. Conclusively, PC represents a promising culture material for stem cells.

## **Acknowledgements**

The authors acknowledge Robert Jeziorski and Mario Rettschlag for preparation of sterilized PS and PC inserts. This work was financially supported by the Helmholtz Association

(Helmholtz-Portfolio Topic “Technology and Medicine”, Helmholtz Virtual Institute “Multifunctional Biomaterials for Medicine” (VH-VI-423), Helmholtz Graduate School of Macromolecular Bioscience (MacroBio), and programme-oriented funding).

## References

- [1] S.T.F. Hsiao, A. Asgari, Z. Lokmic, R. Sinclair, G.J. Disting, S.Y. Lim, R.J. Dilley, Comparative Analysis of Paracrine Factor Expression in Human Adult Mesenchymal Stem Cells Derived from Bone Marrow, Adipose, and Dermal Tissue, *Stem Cells Dev* 21(12) (2012) 2189-2203.
- [2] L. de Girolamo, E. Lucarelli, G. Alessandri, M.A. Avanzini, M.E. Bernardo, E. Biagi, A.T. Brini, G. D'Amico, F. Fagioli, I. Ferrero, F. Locatelli, R. Maccario, M. Marazzi, O. Parolini, A. Pessina, M.L. Torre, I.M.S.C. Grp, Mesenchymal Stem/Stromal Cells: A New "Cells as Drugs" Paradigm. Efficacy and Critical Aspects in Cell Therapy, *Curr Pharm Design*. 19(13) (2013) 2459-2473.
- [3] J.F. Stoltz, D. Bensoussan, N. De Isla, L. Zhang, Z. Han, J. Magdalou, C. Huselstein, J.S. Ye, B. Leballe, V. Decot, L. Reppel, Stem cells and vascular regenerative medicine: A mini review, *Clin Hemorheol Micro* 64(4) (2016) 613-633.
- [4] S. Passeri, F. Nocchi, R. Lamanna, S. Lapi, V. Miragliotta, E. Giannessi, F. Abramo, M.R. Stornelli, M. Matarazzo, D. Plenteda, P. Urciuoli, F. Scatena, A. Coli, Isolation and expansion of equine umbilical cord-derived matrix cells (EUCMCs), *Cell Biol Int* 33(1) (2009) 100-105.
- [5] C.A. Toupadakis, A. Wong, D.C. Genetos, W.K. Cheung, D.L. Borjesson, G.L. Ferraro, L.D. Galuppo, J.K. Leach, S.D. Owens, C.E. Yellowley, Comparison of the osteogenic potential of equine mesenchymal stem cells from bone marrow, adipose tissue, umbilical cord blood, and umbilical cord tissue, *Am J Vet Res* 71(10) (2010) 1237-1245.
- [6] O. Felthaus, T. Schon, D. Schiltz, T. Aung, B. Kuhlmann, F. Jung, A. Anker, S. Klein, L. Prantl, Adipose tissue-derived stem cells from affected and unaffected areas in patients with multiple symmetric lipomatosis show differential regulation of mTOR pathway genes, *Clin Hemorheol Micro* 69(1-2) (2018) 141-151.
- [7] J.C. Estrada, Y. Torres, A. Benguria, A. Dopazo, E. Roche, L. Carrera-Quintanar, R.A. Perez, J.A. Enriquez, R. Torres, J.C. Ramirez, E. Samper, A. Bernad, Human mesenchymal stem cell-replicative senescence and oxidative stress are closely linked to aneuploidy, *Cell Death Dis* 4 (2013) e691.
- [8] F. Gattazzo, A. Urciuolo, P. Bonaldo, Extracellular matrix: A dynamic microenvironment for stem cell niche, *Bba-Gen Subjects* 1840(8) (2014) 2506-2519.
- [9] Z.D. Li, W.W. Wang, X. Xu, K. Kratz, J. Zou, L. Lysyakova, M. Heuchel, A. Kurtz, M. Gossen, N. Ma, A. Lendlein, Integrin beta 1 activation by micro-scale curvature promotes pro-angiogenic secretion of human mesenchymal stem cells, *J Mater Chem B* 5(35) (2017) 7415-7425.
- [10] C.J. Mann, E. Perdiguero, Y. Kharraz, S. Aguilar, P. Pessina, A.L. Serrano, P. Munoz-Canoves, Aberrant repair and fibrosis development in skeletal muscle, *Skelet Muscle* 1 (2011) 1-21.
- [11] T.M. Jiang, G.J. Xu, Q.Y. Wang, L.H. Yang, L. Zheng, J.M. Zhao, X.D. Zhang, In vitro expansion impaired the stemness of early passage mesenchymal stem cells for treatment of cartilage defects, *Cell Death Dis* 8 (2017) 2851-2862.
- [12] W.E. Siebert, S. Mai, S. Kurtz, Retrieval analysis of a polycarbonate-urethane acetabular cup: A case report, *J Long Term Eff Med Implants* 18(1) (2008) 69-74.

- [13] Y.W. Tang, R.S. Labow, I. Revenko, J.P. Santerre, Influence of surface morphology and chemistry on the enzyme catalyzed biodegradation of polycarbonate-urethanes, *J Biomat Sci-Polym E* 13(4) (2002) 463-483.
- [14] G. Humme, H. Rohr, V. Serini, Structure and Properties of Multiphase Plastics .2. Rubber Modified Polycarbonates, *Angew Makromol Chem* 58 (1977) 85-94.
- [15] W. Wang, N. Ma, K. Kratz, X. Xu, Z. Li, T. Roch, K. Bieback, F. Jung, A. Lendlein, The influence of polymer scaffolds on cellular behaviour of bone marrow derived human mesenchymal stem cells, *Clin Hemorheol Microcirc* 52(2-4) (2012) 357-73.
- [16] B. Hiebl, K. Lutzow, M. Lange, F. Jung, B. Seifert, F. Klein, T. Weigel, K. Kratz, A. Lendlein, Cytocompatibility testing of cell culture modules fabricated from specific candidate biomaterials using injection molding, *J Biotechnol* 148(1) (2010) 76-82.
- [17] T. Razafiarison, C.N. Holenstein, T. Stauber, M. Jovic, E. Vertudes, M. Loparic, M. Kawecki, L. Bernard, U. Silvan, J.G. Snedeker, Biomaterial surface energy-driven ligand assembly strongly regulates stem cell mechanosensitivity and fate on very soft substrates, *P Natl Acad Sci USA* 115(18) (2018) 4631-4636.
- [18] B. Cao, Y.M. Peng, X.N. Liu, J.D. Ding, Effects of Functional Groups of Materials on Nonspecific Adhesion and Chondrogenic Induction of Mesenchymal Stem Cells on Free and Micropatterned Surfaces, *Acs Appl Mater Inter* 9(28) (2017) 23574-23585.
- [19] H.W. Lv, L.S. Li, M.Y. Sun, Y. Zhang, L. Chen, Y. Rong, Y.L. Li, Mechanism of regulation of stem cell differentiation by matrix stiffness, *Stem Cell Res Ther* 6 (2015) 103-113.
- [20] K. Metavarayuth, P. Sitasuwan, X. Zhao, Y. Lin, Q. Wang, Influence of Surface Topographical Cues on the Differentiation of Mesenchymal Stem Cells in Vitro, *Acs Biomater Sci Eng* 2(2) (2016) 142-151.
- [21] D.A. Lee, M.M. Knight, J.J. Campbell, D.L. Bader, Stem Cell Mechanobiology, *J Cell Biochem* 112(1) (2011) 1-9.
- [22] R.M. Delaine-Smith, G.C. Reilly, Mesenchymal stem cell responses to mechanical stimuli, *Muscles Ligaments Tendons J* 2(3) (2012) 169-80.
- [23] S.D. Subramony, B.R. Dargis, M. Castillo, E.U. Azeloglu, M.S. Tracey, A. Su, H.H. Lu, The guidance of stem cell differentiation by substrate alignment and mechanical stimulation, *Biomaterials* 34(8) (2013) 1942-53.
- [24] K.E. Michael, D.W. Dumbauld, K.L. Burns, S.K. Hanks, A.J. Garcia, Focal Adhesion Kinase Modulates Cell Adhesion Strengthening via Integrin Activation, *Mol Biol Cell* 20(9) (2009) 2508-2519.
- [25] Y. Sawada, M. Tamada, B.J. Dubin-Thaler, O. Cherniavskaya, R. Sakai, S. Tanaka, M.P. Sheetz, Force sensing by mechanical extension of the Src family kinase substrate p130Cas, *Cell* 127(5) (2006) 1015-1026.
- [26] A.E.X. Brown, D.E. Discher, Conformational Changes and Signaling in Cell and Matrix Physics, *Curr Biol* 19(17) (2009) R781-R789.
- [27] R. Zaidel-Bar, M. Cohen, L. Addadi, B. Geiger, Hierarchical assembly of cell-matrix adhesion complexes, *Biochem Soc T* 32 (2004) 416-420.
- [28] A.J. Engler, S. Sen, H.L. Sweeney, D.E. Discher, Matrix elasticity directs stem cell lineage specification, *Cell* 126(4) (2006) 677-689.
- [29] N.A. Hotchin, A. Hall, The assembly of integrin adhesion complexes requires both extracellular matrix and intracellular rho/rac GTPases, *J Cell Biol* 131(6) (1995) 1857-1865.
- [30] G. Lepperdinger, R. Brunauer, A. Jamnig, G. Laschober, M. Kassem, Controversial issue: Is it safe to employ mesenchymal stem cells in cell-based therapies?, *Exp Gerontol* 43(11) (2008) 1018-1023.
- [31] L. Hayflick, The Limited in Vitro Lifetime of Human Diploid Cell Strains, *Exp Cell Res* 37 (1965) 614-36.

- [32] K. Stenderup, J. Justesen, C. Clausen, M. Kassem, Aging is associated with decreased maximal life span and accelerated senescence of bone marrow stromal cells, *Bone* 33(6) (2003) 919-926.
- [33] N. Gago, V. Perez-Lopez, J.P. Sanz-Jaka, P. Cormenzana, I. Eizaguirre, A. Bernad, A. Izeta, Age-Dependent Depletion of Human Skin-Derived Progenitor Cells, *Stem Cells* 27(5) (2009) 1164-1172.
- [34] A.Y. Maslov, T.A. Barone, R.J. Plunkett, S.C. Pruitt, Neural stem cell detection, characterization, and age-related changes in the subventricular zone of mice, *J Neurosci* 24(7) (2004) 1726-1733.
- [35] A.V. Molofsky, S.G. Slutsky, N.M. Joseph, S. He, R. Pardal, J. Krishnamurthy, N.E. Sharpless, S.J. Morrison, Increasing p16INK4a expression decreases forebrain progenitors and neurogenesis during ageing, *Nature* 443(7110) (2006) 448-52.
- [36] D.L. Jones, T.A. Rando, Emerging models and paradigms for stem cell ageing, *Nat Cell Biol* 13(5) (2011) 506-12.
- [37] V. Turinetti, E. Vitale, C. Giachino, Senescence in Human Mesenchymal Stem Cells: Functional Changes and Implications in Stem Cell-Based Therapy, *Int J Mol Sci* 17(7) (2016) 1164-1181.
- [38] F. Rodier, D.P. Munoz, R. Teachenor, V. Chu, O. Le, D. Bhaumik, J.P. Coppe, E. Campeau, C.M. Beausejour, S.H. Kim, A.R. Davalos, J. Campisi, DNA-SCARS: Distinct nuclear structures that sustain damage-induced senescence growth arrest and inflammatory cytokine secretion, *J Cell Sci* 124(1) (2011) 68-81.
- [39] A. Banfi, A. Muraglia, B. Dozin, M. Mastrogiacomo, R. Cancedda, R. Quarto, Proliferation kinetics and differentiation potential of ex vivo expanded human bone marrow stromal cells: Implications for their use in cell therapy, *Exp Hematol* 28(6) (2000) 707-715.
- [40] S. Geissler, M. Textor, J. Kuhnisch, D. Konnig, O. Klein, A. Ode, T. Pfitzner, J. Adjaye, G. Kasper, G.N. Duda, Functional Comparison of Chronological and In Vitro Aging: Differential Role of the Cytoskeleton and Mitochondria in Mesenchymal Stromal Cells, *Plos One* 7(12) (2012) 52700-52712.
- [41] K. Kornicka, K. Marycz, K.A. Tomaszewski, M. Maredziak, A. Smieszek, The Effect of Age on Osteogenic and Adipogenic Differentiation Potential of Human Adipose Derived Stromal Stem Cells (hASCs) and the Impact of Stress Factors in the Course of the Differentiation Process, *Oxid Med Cell Longev* 2015 (2015) 309169-309188.
- [42] S. Geissler, M. Textor, J. Kuhnisch, D. Konnig, O. Klein, A. Ode, T. Pfitzner, J. Adjaye, G. Kasper, G.N. Duda, Functional comparison of chronological and in vitro aging: differential role of the cytoskeleton and mitochondria in mesenchymal stromal cells, *Plos One* 7(12) (2012) e52700.
- [43] V.B.R. Konala, M.K. Mamidi, R. Bhonde, A.K. Das, R. Pochampally, R. Pal, The current landscape of the mesenchymal stromal cell secretome: A new paradigm for cell-free regeneration, *Cytotherapy* 18(1) (2016) 13-24.
- [44] L.D. Meirelles, A.M. Fontes, D.T. Covas, A.I. Caplan, Mechanisms involved in the therapeutic properties of mesenchymal stem cells, *Cytokine Growth F R* 20(5-6) (2009) 419-427.
- [45] W.S. Park, S.Y. Ahn, S.I. Sung, J.Y. Ahn, Y.S. Chang, Strategies to enhance paracrine potency of transplanted mesenchymal stem cells in intractable neonatal disorders, *Pediatr Res* 83(1) (2018) 214-222.
- [46] A. Hoeben, B. Landuyt, M.S. Highley, H. Wildiers, A.T. Van Oosterom, E.A. De Bruijn, Vascular endothelial growth factor and angiogenesis, *Pharmacol Rev* 56(4) (2004) 549-580.