



Final Draft **of the original manuscript**

Fehn, N.; Kronbichler, M.; Munch, P.; Wall, W.:

Numerical evidence of anomalous energy dissipation in incompressible Euler flows: towards grid-converged results for the inviscid Taylor-Green problem.

In: Journal of Fluid Mechanics. Vol. 932 (2022) A40.

First published online by Cambridge University Press: 15.12.2021

<https://dx.doi.org/10.1017/jfm.2021.1003>

Numerical evidence of anomalous energy dissipation in incompressible Euler flows: Towards grid-converged results for the inviscid Taylor–Green problem

5 **Niklas Fehn¹†, Martin Kronbichler¹, Peter Munch^{1,2} and Wolfgang A. Wall¹**

¹Institute for Computational Mechanics, Technical University of Munich, Boltzmannstr. 15, 85748 Garching, Germany

²Institute of Material Systems Modeling, Helmholtz-Zentrum Hereon, Max-Planck-Str. 1, 21502 Geesthacht, Germany

The well-known energy dissipation anomaly in the inviscid limit, related to velocity singularities according to Onsager, still needs to be demonstrated by numerical experiments. The present work contributes to this topic by high-resolution numerical simulations of the inviscid three-dimensional Taylor–Green vortex problem using a novel high-order discontinuous Galerkin discretisation approach for the incompressible Euler equations. The main methodological ingredient is the use of a discretisation scheme with inbuilt dissipation mechanisms as opposed to discretely energy-conserving schemes, which – by construction – rule out the occurrence of anomalous dissipation. We investigate effective spatial resolution up to 8192^3 (defined based on the 2π -periodic box) and make the interesting phenomenological observation that the kinetic energy evolution does not tend towards exact energy conservation for increasing spatial resolution of the numerical scheme, but that the sequence of discrete solutions seemingly converges to a solution with nonzero kinetic energy dissipation rate. By taking the fine-resolution simulation as reference, we measure grid-convergence with a relative L^2 -error of 0.27% for the temporal evolution of the kinetic energy and 3.52% for the kinetic energy dissipation rate against the dissipative fine-resolution simulation. The present work raises the question whether such results can be seen as a numerical confirmation of the famous energy dissipation anomaly. Due to the relation between anomalous energy dissipation and the occurrence of singularities for the incompressible Euler equations according to Onsager’s conjecture, we elaborate on an indirect approach for the identification of finite-time singularities that relies on energy arguments.

30 **Key words:** anomalous energy dissipation, finite-time singularities, incompressible Euler equations, Onsager’s conjecture, turbulence

1. Motivation

Singularities play a key role in fluid mechanics (Eggers 2018). While singularities in the form of shocks are well-understood for the compressible Euler equations and the inviscid Burgers equation (Burgers 1948) as a simplified model, the occurrence of singularities that develop in finite time is discussed controversially for the three-dimensional incompressible Euler equations.

† Email address for correspondence: niklas.fehn@tum.de

The occurrence of singularities is strongly related to anomalous dissipation of kinetic energy in three-dimensional incompressible Euler flows according to the pioneering work by Onsager (1949), which is well-documented in review articles (Eyink & Sreenivasan 2006; Eyink 2008) and in the recent essay by Dubrulle (2019). Due to the relation between singularities and dissipation, we distinguish between (i) a *direct* approach to identify finite-time singularities for incompressible Euler flows, e.g., by showing that the vorticity blows up in finite time through different methods (e.g., an analysis of the kind $\|\omega\|_\infty \sim (t_* - t)^{-\gamma}$ according to the Beale–Kato–Majda theorem (Beale *et al.* 1984) trying to identify t_* and γ from numerical results), and (ii) an *indirect* approach providing indications of finite-time singularities by observing an “anomalous” dissipative behaviour in the kinetic energy evolution. While most approaches in the literature can be identified as belonging to the first category, the present work focuses on a technique related to the second category. As explained below, this *indirect* approach is rather new or unexplored due to certain subtleties with respect to numerical discretisation schemes. To complement these results, we additionally show numerical results related to the direct identification approach such as the temporal evolution of the maximum vorticity $\|\omega\|_\infty$ and the enstrophy \mathcal{E} .

1.1. *State-of-the-art and limitations in tracing finite-time singularities*

A strategy to identify potential singularities directly are time series expansions (Brachet *et al.* 1983; Pelz & Gulak 1997; Taylor & Green 1937; Morf *et al.* 1980), but it was found that numerical inaccuracies prevent a definite answer when using this technique. Numerical investigations by means of PDE (partial differential equation) solvers have therefore played the most dominant role in the exploration of finite-time singularities as detailed below.

A difficulty in identifying singularities with the direct approach by numerical simulations is the inherent conflict that arbitrarily small structures can not be resolved with a numerical simulation of finite resolution, which renders this problem one of the most challenging topics in computational fluid dynamics. Much work has been done in this field. In the 1980s and 1990s, several early works on direct numerical simulation (DNS) of both inviscid and high-Reynolds-number viscous incompressible flows reported indications of finite-time singularities for the incompressible Euler equations (Brachet *et al.* 1983; Kerr & Hussain 1989; Kerr 1993; Brachet *et al.* 1992; Boratav & Pelz 1994). Symmetry in the initial conditions plays an important role as specifically mentioned and addressed by some works (Boratav & Pelz 1994; Pelz & Gulak 1997; Pelz 2001), raising the question whether singularities are possible for problems that are not perfectly symmetric. However, these works do not allow a definite answer to the question of finite-time singularities, see also the review articles on this topic (Gibbon 2008; Hou & Li 2008). One of the main reasons why the results of these studies have been inconclusive is that the spatial resolution has been limited due to the computational power and computational approaches available at the time. Numerical results shown in Hou & Li (2008) suggest that dynamic depletion of vortex stretching could be a mechanism that prevents a finite-time blow-up, but the same authors report evidence for a finite-time singularity for a different flow configuration with solid boundaries in a later work (Luo & Hou 2014).

In terms of the flow configuration being studied, numerical investigations on finite-time singularities can be categorised as follows: The Taylor–Green vortex (TGV) has been analyzed in Brachet *et al.* (1983, 1992); Brachet (1991); Shu *et al.* (2005); Cichowlas & Brachet (2005); Bustamante & Brachet (2012) and for a regularised problem considering the Euler–Voigt equations in Larios *et al.* (2018), the high-symmetry Kida–Pelz initial condition in Hou & Li (2008); Cichowlas & Brachet (2005); Grafke *et al.* (2008), colliding Lamb dipoles in Orlandi *et al.* (2012), and other perturbed cylindrical vortex tubes in Kerr & Hussain (1989); Kerr (1993); Hou & Li (2008); Grauer *et al.* (1998); Kerr (2013). Most studies used spectral methods as discretisation schemes.

For these direct numerical simulations, common approaches to trace singularities are moni-

toring the maximum vorticity $\|\omega\|_\infty$ over time, see the Beale–Kato–Majda theorem (Beale *et al.* 1984), and the “analyticity strip” method (see Sulem *et al.* 1983), which aims at capturing the smallest scales of the flow. The width of the analyticity strip $\delta(t)$, obtained from fitting the energy spectrum to $E(k, t) = C(t)k^{-n(t)} \exp(-2k\delta(t))$, is monitored over time for successively
 90 finer spatial resolutions up to a resolution for which extrapolations of $\delta(t)$ allow conclusions whether $\delta(t)$ reaches 0 in a finite time (finite-time singularity) or whether $\delta(t)$ decreases only exponentially in time (regularity at all times).

Numerical results for the three-dimensional inviscid Taylor–Green vortex shown in Brachet
et al. (1983); Cichowlas & Brachet (2005) indicate only an exponential decay, but this might be
 95 due to the limited spatial resolution and also due to the fact that only small times of the TGV flow have been considered, so that a finite-time singularity can not be excluded from these results. In a later work by Bustamante & Brachet (2012), a change in regime indicating potentially faster than exponential decay is reported and the results are “not inconsistent with the occurrence of a singularity”, but again resolutions higher than the maximum one of 4096^3 would be required for
 100 definite answers. In Cichowlas & Brachet (2005) it is estimated that conclusions regarding finite-time singularities using the analyticity strip method would require spatial resolutions of $(16k)^3$ to $(32k)^3$ for the Kida–Pelz initial data. A recent study by Campolina & Mailybaev (2018) suggests that the resolution available via classical DNS is not sufficient to investigate blow-up.

The development of pancake-like structures with exponentially growing vorticity during the
 105 early development of turbulence from smooth initial data is studied in Agafontsev *et al.* (2015). In Kerr (2013), a new kind of analysis based on rescaled vorticity moments is proposed studying anti-parallel vortex tubes, and only double-exponential growth in vorticity is observed as opposed to the singular behaviour suspected in a previous work by Kerr (1993). A model describing a cascade of transformations between vortex filaments and sheets potentially explaining the
 110 mechanism of singularity formation in the Euler equations is proposed in Brenner *et al.* (2016). Another model has been described recently in Moffatt (2019). In McKeown *et al.* (2018), an iterative cascade of instabilities for head-on collisions of vortex rings is investigated both experimentally and numerically.

1.2. Energy dissipation anomaly

We now focus on the evolution of kinetic energy in incompressible Euler flows. Of particular interest is the question whether inviscid flows are able to dissipate energy, and if so, by which mechanism such a behaviour can be explained, given that no viscous effects are present. The kinetic energy dissipation equation – valid for incompressible viscous ($\nu > 0$) flows with continuously differentiable solution on a domain with periodic boundaries – reads (Onsager 1949; Eyink & Sreenivasan 2006)

$$\frac{\partial E(t, \nu)}{\partial t} = - \int_{\Omega} \nu \nabla \mathbf{u}^\nu : \nabla \mathbf{u}^\nu \, d\Omega,$$

which implies conservation of energy in the inviscid limit $\nu = 0$ provided that the solution is sufficiently regular. However, from phenomenological descriptions of turbulence, there is empirical evidence that the dissipation rate does not tend to zero in the limit $\text{Re} \rightarrow \infty$ or $\nu \rightarrow 0$ but takes a positive value independent of ν , which is known as dissipation anomaly or the zeroth law of turbulence (Eyink 2008; Dubrulle 2019). As noted in Eyink (2008), this has first been observed by Taylor (1935), and also Kolmogorov’s similarity theory of turbulence (Kolmogorov 1991) is based on the assumption of a non-vanishing energy dissipation rate in the inviscid limit. Numerical evidence that the dissipation rate is independent of ν for large Re is for example given in Sreenivasan (1998); Kaneda *et al.* (2003); Orlandi *et al.* (2012), and experimental evidence for example in Pearson *et al.* (2002); Dubrulle (2019). Under certain regularity or smoothness assumptions (existence of a strong L^3 -limit; we refer to Duchon & Robert (2000); Drivas &

Nguyen (2019); Drivas & Eyink (2019) for a precise discussion), weak Euler solutions are the $\nu \rightarrow 0$ limit of Leray–Hopf weak solutions \mathbf{u}^ν of the Navier–Stokes equations, so that the dissipation rate in the inviscid limit equals the viscous dissipation rate in the limit $\nu \rightarrow 0$

$$\frac{\partial E_{\nu=0}(t)}{\partial t} = \lim_{\nu \rightarrow 0} \frac{\partial E(t, \nu)}{\partial t} = \lim_{\nu \rightarrow 0} -\nu \int_{\Omega} \nabla \mathbf{u}^\nu : \nabla \mathbf{u}^\nu \, d\Omega = -D(t) \leq 0, \quad (1.1)$$

115 where anomalous energy dissipation means that $D(t) > 0$ for some (or all) $t > t_*$.[†] In general, weak Euler solutions may neither be unique nor the zero-viscosity-limit of weak Navier–Stokes solutions, which might themselves be non-unique (Isett 2017; Buckmaster & Vicol 2019, 2020; Daneri *et al.* 2021; Buckmaster *et al.* 2021). In this sense, we note that the first equality in equation (1.1) is a conditional one. Let us also refer to Brenier *et al.* (2011); Wiedemann (2017)
120 regarding the topic of weak–strong uniqueness of Euler solutions.

The above argumentation already indicates that the theory explaining dissipation of energy in the absence of viscosity is related to the spatial regularity of the solution. According to Onsager (1949), energy dissipation in three-dimensional incompressible flows can take place in the absence of viscosity by the formation of singularities,[‡] with the cascade from large to (arbitrarily) small
125 scales taking place in finite time, see also Eyink & Sreenivasan (2006). According to Onsager’s conjecture, dissipation of energy may occur if the velocity is Hölder continuous with exponent $\leq 1/3$ (while Onsager’s assertion says that energy is conserved for exponents $> 1/3$, see Eyink (1994); Constantin *et al.* (1994); Duchon & Robert (2000); Cheskidov *et al.* (2008) for proofs). For mathematical literature dealing with proofs of Onsager’s conjecture we refer to De Lellis
130 & Székelyhidi Jr. (2013, 2014); Buckmaster *et al.* (2016, 2018); Isett (2018) and references therein, where dissipative weak Euler solutions up to Onsager’s critical regularity have been constructed using convex integration techniques. Newest insights from these works thus confirm that the Hölder exponent of $1/3$ is indeed the critical one in terms of energy dissipation. As noted in Dubrulle (2019), the original Kolmogorov cascade picture implies irregularities of the velocity
135 field (at least locally) with Hölder exponent $\leq 1/3$, but it was Onsager who established the link between energy dissipation and irregularities of the velocity field for the Euler equations.

The above considerations might explain why this phenomenon is denoted as “kinetic energy dissipation anomaly”, an alternative term used e.g. in Dubrulle (2019) is “inertial dissipation” (as opposed to viscous dissipation). The one-dimensional inviscid Burgers equation (Burgers
140 1948) with formation of a shock and the associated dissipation of energy serves as a prominent and well-understood example, and is for example discussed in Sulem *et al.* (1983) in the context of finite-time singularities and in Dubrulle (2019) in the context of inertial energy dissipation. In Josserand *et al.* (2020), the phenomenon of energy dissipation through finite-time singularities is illustrated for another one-dimensional model problem, the nonlinear Schrödinger equation. For
145 the two-dimensional incompressible Euler equations, it is known that singularities can not develop in finite time from smooth initial data (Eyink & Sreenivasan 2006). Regarding three-dimensional turbulent flows, Onsager’s conjecture appears to be widely accepted by now with the occurrence of singularities representing a building block of modern understandings of turbulence (Dubrulle 2019).

[†] As a consequence, the enstrophy is inversely proportional to the viscosity for large Reynolds numbers in the case of anomalous dissipation with $D(t) > 0$.

[‡] Hence, the mechanism explaining the occurrence of kinetic energy dissipation in the limit $\nu \rightarrow 0$ is that the velocity gradient might tend to infinity in this limit. Literally, Onsager (1949) wrote: “In the absence of viscosity, the standard proof of the conservation of energy does not apply, because the velocity field does not remain differentiable!”. Interestingly, Onsager did not consider the energy-dissipating behaviour of inviscid flows (“ideal turbulence”) an anomalous behaviour but rather a matter of fact.

1.3. Interplay between physics and numerics

The present work focuses on the numerical solution of turbulent flows and particularly the inviscid limit. Having a closer look at numerical simulations of the inviscid Taylor–Green vortex, it can be observed that many of these simulations have been performed mainly for small times up to $t \approx 5$ (up to $t = 4$ in Brachet *et al.* (1983, 1992); Cichowlas & Brachet (2005); Bustamante & Brachet (2012) and up to $t \leq 6$ in Shu *et al.* (2005); Chapelier *et al.* (2012)), but not beyond the time at which finite-time singularities have been suspected, especially not up to the time at which the transition to a fully turbulent state takes place, with maximum kinetic energy dissipation rate at time $t = 8–9$ (expected from high-Reynolds number viscous simulations (see Brachet *et al.* 1983)) and subsequent decaying turbulence. As mentioned by some of these works, one reason for this is that the results for a specific resolution are no longer reliable at later times once the flow becomes under-resolved (note that the spatial resolution is severely limited by computational resources even for large supercomputers available today). The key aspect, however, is that numerical simulations of the incompressible Euler equations are very challenging in terms of energy stability and numerical blow-up of the discretisation scheme. Often, a lack of robustness of the numerical discretisation scheme is reported for this challenging inviscid Taylor–Green vortex problem (see for example Chapelier *et al.* 2012; Winters *et al.* 2018).

The lack of understanding of what is to be expected in terms of kinetic energy dissipation from a physical perspective (energy dissipation anomaly discussed in section 1.2) manifests itself in an uncertainty regarding the optimal design of discretisation schemes from a numerical perspective. In Moura *et al.* (2017*b,a*); Piatkowski (2019), it is argued that a fundamentally different behaviour in terms of energy dissipation and time reversibility is expected between viscous flows in the limit $\text{Re} \rightarrow \infty$ and inviscid flows at $\text{Re} = \infty$. Especially in numerical studies, it is often assumed that energy conservation holds for an exact solution of the Euler equations not only in two space dimensions but also in three space dimensions, see for example Shu *et al.* (2005); Bustamante & Brachet (2012); Grauer *et al.* (1998); Chapelier *et al.* (2012); Winters *et al.* (2018); Schroeder (2019); Kraus *et al.* (2020) and the recent review article by Coppola *et al.* (2019) to mention just a few. Numerical schemes that are exactly energy-conserving can indeed be constructed and have the advantage that non-linear blow-up of the numerical discretisation scheme can be avoided in the challenging inviscid limit. For these reasons, energy-conserving schemes appear to be the current gold standard for the simulation of this type of problems. Inviscid TGV simulations performed in Schroeder (2019) using exactly divergence-free, energy-conserving discretisation methods result in an exact conservation of energy, and the results are considered superior as compared to simulations with upwind fluxes that show a dissipative behaviour.

However, the use of energy-conserving schemes is accompanied by a major limitation, namely that it excludes – by construction – the occurrence of anomalous energy dissipation. Onsager’s conjecture dictates to rethink whether it is really a desirable quality criterion that a numerical method preserves the kinetic energy exactly in the inviscid limit $\nu = 0$. If Onsager’s hypothesis is true, there is an inconsistency between the physical dissipation behaviour and the numerical dissipation behaviour of energy-conserving discretisation methods. Then, energy-conserving numerical methods would result in an $O(1)$ error in the case of inviscid flows with anomalous/inertial energy dissipation. Since no energy can leave the system for such a discretisation scheme, energy accumulates in small scales, a well-known phenomenon called thermalisation. In terms of the kinetic energy spectrum, an energy-conserving numerical scheme typically leads to an unphysical equipartitioning of energy when simulating beyond the time of the finite-time singularity (Ray *et al.* 2011; Orlandi 2009; Orlandi *et al.* 2012). The energy-conserving results of Schroeder (2019) indeed show such a behaviour. One may conclude that the application of energy-conserving numerical methods is only reasonable for times $t < t_*$ before a potential singularity forms, since anomalous dissipation might occur afterwards.

Further, one may formulate that a numerical scheme must contain mechanisms of dissipation as a minimal requirement in order to address the topic of anomalous energy dissipation. To describe discretisation schemes suitable for investigating anomalous dissipation more precisely, we consider it a prerequisite to use consistent and stable discretisation schemes whose dissipation mechanisms are coupled to under-resolution effects in the numerical approximation of the solution, i.e., the dissipation mechanisms act on the finest resolved scales and shift to smaller scales under mesh refinement. As we discuss in more detail below, it appears to be unclear mathematically whether such a scheme is able to find a dissipative weak solution of the Euler equations.

By studying Galerkin-truncated, energy-conserving simulations of the one-dimensional Burgers equation, the work by Ray *et al.* (2011) describes an interesting phenomenon, called “tyger phenomenon” in that work, where short-wavelength oscillations occur out-of-the-blue in the presence of singularities, that finally lead to thermalisation. The importance of numerical dissipation to avoid the effect of thermalisation for this one-dimensional Burgers problem is emphasised in a recent study by Murugan *et al.* (2020), which can therefore be seen in close analogy to the present work focusing on 3D Euler.

The situation is less complicated for two-dimensional Euler flows that are non-dissipative. In that case, it can be expected that the kinetic energy dissipation rate converges to zero under mesh refinement for a consistent and energy-stable discretisation scheme, and that there is per se no conflict with physics if an energy-conserving scheme is applied.

1.4. An indirect approach to identify finite-time singularities by energy considerations

The *indirect* approach to identify finite-time singularities relies on the physical intuition that the appearance of anomalous energy dissipation in free decay from smooth initial data requires a finite-time singularity. The basic idea is to capture the temporal evolution of the kinetic energy by a numerical method with appropriate inbuilt dissipation mechanisms as described in Section 1.3. If grid-convergence to a dissipative solution with non-zero kinetic energy dissipation rate can be demonstrated numerically, indirect evidence of a finite-time singularity is provided by the following line of arguments:

- (i) Assume convergence of a sequence of numerical solutions to a dissipative weak Euler solution for $h \rightarrow 0$.
- (ii) Weak–strong uniqueness holds for dissipative weak Euler solutions (Lions 1996; Brenier *et al.* 2011; Wiedemann 2017).
- (iii) Supposing that an energy dissipation anomaly with non-zero kinetic energy dissipation rate is observed, it follows from items **i** and **ii** that a strong solution cannot exist but must have become singular.

The conclusion in item **iii** is based on an indirect proof. Assume that a strong solution exists. This strong solution is energy-conserving. By weak–strong uniqueness, the weak solution must be identical and, therefore, energy-conserving. By contradiction it follows that a strong solution cannot exist.

Note that assumption **i** could be weakened. Rather than **i**, only convergence to a dissipative generalised weak solution in the sense of Lions (1996) or DiPerna & Majda (1987) is required. Interestingly, this weaker assumption might be provable for the limit $h \rightarrow 0$, as it has in fact been proven for the limit $\nu \rightarrow 0$ (at least along sub-sequences). Since the existence of generalised weak Euler solutions as limits along sub-sequences relies on very general compactness arguments that require only L^2 (kinetic energy) bounds for both the Lions and DiPerna & Majda theories, a proof for the limit $h \rightarrow 0$ for a DG Euler discretisation scheme as used here might be conceivable in analogy to what has been shown for the limit $\nu \rightarrow 0$. To the best of our knowledge, such a proof is still outstanding.




Figure 1: Temporal evolution of kinetic energy and kinetic energy dissipation rate for the three-dimensional Taylor–Green vortex problem for increasing Reynolds number of $Re = 100, 200, 400, 800, 1600, 3200, \infty$. For each Re , results are shown for two mesh resolutions (fine mesh as solid line, coarse mesh as dashed-dotted line). The effective resolutions (see section 3 for a definition) are $64^3, 128^3$ for $Re = 100$, $128^3, 256^3$ for $Re = 200, 400$, $256^3, 512^3$ for $Re = 800$, $1024^3, 2048^3$ for $Re = 1600$, $2048^3, 4096^3$ for $Re = 3200$, and $4096^3, 8192^3$ for $Re = \infty$. The results suggest that the kinetic energy reduces to a value as low as approximately 0.02 at time $t = 20$ for large Reynolds numbers, and that a similar amount of energy dissipation takes place also in the inviscid limit.

Let us note that a similar idea to identify singularities experimentally based on energy arguments has been used in Saw *et al.* (2016); Kuzzay *et al.* (2017) by calculating the inertial dissipation at scale l from PIV (particle image velocimetry) measurements. To the best of the authors' knowledge, the present study makes first attempts in using energy arguments for singularity detection in numerical simulations of three-dimensional Euler flows.

From numerical simulations of viscous problems at finite Reynolds number, there are indications that the zeroth law of turbulence holds for the Taylor–Green vortex problem. Numerical results for the kinetic energy dissipation rate for increasing Reynolds numbers up to $Re = 3000$ shown in Brachet *et al.* (1983), additional results for a higher Reynolds number of $Re = 5000$ shown in Brachet (1991), $Re = 10000$ in Arndt *et al.* (2020), and $Re = 20000$ in Lamballais *et al.* (2019) strongly suggest that the function $D(t)$ does not tend to zero in the limit $\nu \rightarrow 0$. This argument is summarised in figure 1 showing results for viscous and inviscid simulations of the TGV obtained with the present discretisation approach. For each Reynolds number, results are shown for two resolutions of the numerical discretisation approach to judge whether the results are mesh-independent. We achieve grid-converged results for all finite Reynolds numbers shown in figure 1. In the inviscid limit, the temporal evolution of the kinetic energy is almost indistinguishable for the two finest resolutions, while small differences are still visible in the energy dissipation rate that slightly differs between the two resolutions 4096^3 and 8192^3 at later times around the dissipation maximum and beyond. However, the onset of dissipation around $t \approx 6$ appears to be grid-converged also for this challenging inviscid simulation. These results are consistent with grid-convergence to a dissipative solution of the incompressible Euler equations for the three-dimensional Taylor–Green problem. Figure 1 therefore summarizes the main result of the present work.

While the accumulation of energy in small scales in case of energy-preserving schemes is unphysical, it can be exploited under certain circumstances in order to gain insights into the physical dissipation behaviour. In Cichowlas *et al.* (2005), an effective dissipation is estimated from the small-scale thermalised energy of energy-conserving, spectrally truncated Euler simulations and it is found that the large-scale Euler dynamics are similar to high-Reynolds

number Navier–Stokes dynamics. Although the underlying numerical methods in that work are different from the present study,[†] there are interesting parallels. The onset of dissipation around $t = 5 - 6$ and the dissipation maximum around $t = 8 - 9$ appear to be very similar to the present results, so that both studies can be seen to complement and verify each other. From this perspective, the study by Cichowlas *et al.* (2005) supports the main conclusions of our work.

Instead of considering a two-parameter limit problem $h \rightarrow 0, \nu \rightarrow 0$ as illustrated in figure 1, the remainder of this work focuses on the one-parameter limit $h \rightarrow 0$ for the inviscid limit $\nu = 0$. Let us explain this decision in more detail. A two-parameter study $h \rightarrow 0, \nu \rightarrow 0$ would technically not be realizable due to the large amount of computational costs required for such simulations: Already for moderate Reynolds numbers of $\text{Re} = O(10^4)$ as considered for example in Arndt *et al.* (2020); Lamballais *et al.* (2019), the spatial resolutions required for grid-convergence are comparable to the highest resolution simulations that we are able to realise in the present work for the inviscid limit. These highest resolution simulations require computational costs of tens of millions of CPUh in the present study, despite the fact that we use a highly efficient implementation that is well-optimized for the hardware under consideration. With the goal to realise spatial resolutions as high as those shown in the present study, computational costs would allow us to only consider a single finite Reynolds number beyond what is shown in figure 1, explaining why we immediately address the inviscid limit $\nu = 0$.

1.5. Outline

The rest of this article is organised as follows. We describe the mathematical model of the incompressible Navier–Stokes or Euler equations and its numerical discretisation in space and time in section 2. Section 3 shows results for the one-dimensional Burgers equation with formation of a shock, a two-dimensional shear layer problem with a numerical investigation of the kinetic energy dissipation in the limit $\nu \rightarrow 0$, and finally the three-dimensional Taylor–Green vortex problem that has been suspected to exhibit finite-time singularities in the inviscid limit. In section 4, we summarise our results, draw conclusions, and raise questions based on the present observations.

2. Numerical methods

We seek numerical solutions to the incompressible Euler equations solved on a domain $\Omega \subset \mathbb{R}^d$ in $d = 2, 3$ space dimensions. These have their origin in the equations for viscous fluids with kinematic viscosity ν described by the incompressible Navier–Stokes equations

$$\frac{\partial \mathbf{u}}{\partial t} + \nabla \cdot (\mathbf{u} \otimes \mathbf{u}) - \nu \nabla^2 \mathbf{u} + \nabla p = \mathbf{0}, \quad (2.1)$$

$$\nabla \cdot \mathbf{u} = 0, \quad (2.2)$$

where \mathbf{u} denotes the d -dimensional velocity vector and p the kinematic pressure. The Euler equations are recovered by setting $\nu = 0$. This system of partial differential equations does not extend to $d = 1$ in a meaningful way, since the incompressibility constraint $\partial u / \partial x = 0$ would imply $u = \text{const.}$ [‡] Let us note that the temporal and spatial discretisation schemes discussed below are generic and hold for both two- and three-dimensional problems, but that there are major differences in terms of the flow physics and the mechanisms that make up the nature of turbulence such as the energy transfer to small scales according to a turbulence cascade in three

[†] While the energy dissipation rate is derived by a postprocessing of results in Cichowlas *et al.* (2005), it is simulated directly in the present work.

[‡] However, the one-dimensional Burgers equation serves as a simplified mathematical model for more complex higher-dimensional problems. While results for the inviscid Burgers equation are presented in section 3, this section deals with discretisations of the incompressible Navier–Stokes equations for $d = 2, 3$.

dimensions (Onsager 1949). This fundamentally different behaviour is attributed to the vortex stretching term in the vorticity form of the Euler equations (Gibbon 2008)

$$\frac{D\boldsymbol{\omega}}{Dt} = (\boldsymbol{\omega} \cdot \nabla) \mathbf{u},$$

where the vortex stretching term on the right-hand side vanishes in two dimensions since the vorticity $\boldsymbol{\omega}$ is perpendicular to the velocity \mathbf{u} in that case.

305 The following two subsections detail the temporal and spatial discretisation of the incompressible Navier–Stokes equations (2.1) and (2.2). Discretisation in time is based on projection methods that solve for velocity and pressure unknowns in different sub-steps of a time step. Discretisation in space is based on high-order discontinuous Galerkin (DG) methods with suitable stabilisation techniques that render the method robust for under-resolved, high-Reynolds number flows.

310 2.1. Temporal discretisation – high-order projection method

Discretisation in time is based on projection methods which aim at obtaining computationally efficient incompressible flow solvers by decoupling the velocity and pressure unknowns (Karniadakis & Sherwin 2005). A convection–diffusion type problem is solved for the velocity and a Poisson equation for the pressure with a subsequent projection of the velocity onto the space of solenoidal vector fields according to the Helmholtz decomposition. We use the high-order dual splitting scheme proposed in Karniadakis *et al.* (1991) which consists of the following four sub-steps

$$\frac{\gamma_0^n \hat{\mathbf{u}} - \sum_{i=0}^{J-1} \alpha_i^n \mathbf{u}^{n-i}}{\Delta t_n} = - \sum_{i=0}^{J-1} \beta_i^n \nabla \cdot (\mathbf{u}^{n-i} \otimes \mathbf{u}^{n-i}), \quad (2.3)$$

$$-\nabla^2 p^{n+1} = - \frac{\gamma_0^n}{\Delta t_n} \nabla \cdot \hat{\mathbf{u}}, \quad (2.4)$$

$$\hat{\mathbf{u}} = \mathbf{u} - \frac{\Delta t_n}{\gamma_0^n} \nabla p^{n+1}, \quad (2.5)$$

$$\frac{\gamma_0^n}{\Delta t_n} \mathbf{u}^{n+1} - \nu \nabla^2 \mathbf{u}^{n+1} = \frac{\gamma_0^n}{\Delta t_n} \hat{\mathbf{u}}, \quad (2.6)$$

where n denotes the current time step in which the equations are integrated from time t_n to time $t_{n+1} = t_n + \Delta t_n$. The temporal discretisation is based on a BDF scheme of order J with coefficients γ_0^n and $\alpha_i^n, i = 0, \dots, J - 1$. In the first sub-step, equation (2.3), the convective term is treated explicitly in time by using a high-order extrapolation scheme of order J with coefficients $\beta_i^n, i = 0, \dots, J - 1$. In the next two sub-steps, a pressure Poisson equation, equation (2.4), is solved and a divergence-free velocity is obtained in the projection step, equation (2.5). Finally, the viscous term is taken into account in the last sub-step, equation (2.6), which can be omitted in case of the Euler equations, $\mathbf{u}^{n+1} = \hat{\mathbf{u}}$. The explicit treatment of the convective term implies a restriction of the time step sizes according to the Courant–Friedrichs–Lewy (CFL) condition. For reasons of computational efficiency, we use adaptive time stepping with variable time step sizes Δt_n by readjusting the time step size after each time step in a way to maximise the time step size and operate close to the CFL stability limit. We introduce the CFL condition below since it also depends on the spatial discretisation scheme. A second-order accurate time integration scheme with $J = 2$ is used in the present work.

325 2.2. Spatial discretisation – high-order discontinuous Galerkin method

A key element of the present study is the use of a novel high-order discontinuous Galerkin discretisation for incompressible flows. This discretisation approach has been developed recently

10

and is documented in a series of publications (Krank *et al.* 2017; Fehn *et al.* 2017, 2018b, 2019), with a focus on the stability of projection methods in Fehn *et al.* (2017), and a focus
 330 on the stability and dissipation characteristics of discontinuous Galerkin discretisations for under-resolved turbulence in Fehn *et al.* (2018b, 2019). Of particular importance are the inbuilt dissipation mechanisms of discontinuous Galerkin methods acting on the finest resolved scales of the flow. In order to investigate the phenomenon of anomalous energy dissipation, we consider it a prerequisite to use a discretisation scheme that provides the flexibility to find both dissipative and
 335 non-dissipative solutions. Let us point out that all simulations shown in this work are performed as direct numerical simulations, i.e., without any form of explicit turbulence model or numerical viscosity. More precisely, upwind-type numerical fluxes represent the main dissipation mechanism of the present scheme, see also Ainsworth (2004) for a general dispersion and dissipation analysis and distribution of dissipation among the resolved scales. Note that the $H(\text{div})$ -stabilization presented below also contributes with a certain amount of dissipation. For reasons of brevity and
 340 to focus on the main aspects, we avoid technical aspects related to the imposition of boundary conditions in the following, but refer to the original publications.

We assume a computational domain $\Omega_h = \bigcup_{e=1}^{N_{\text{el}}} \Omega_e \in \mathbb{R}^d$ consisting of conforming quadrilateral or hexahedral elements Ω_e , $e = 1, \dots, N_{\text{el}}$. A common abstraction of finite element methods is to define a mapping $\mathbf{x}^e(\boldsymbol{\xi})$ from a reference element $\tilde{\Omega}_e = [0, 1]^d$ with Cartesian coordinates $\boldsymbol{\xi}$ to element Ω_e in physical space with coordinates \mathbf{x} , and approximate the solution within each element by polynomials defined in reference coordinates. Here, the numerical solution in $d = 2, 3$ is represented by a tensor product of one-dimensional Lagrange polynomials with a Legendre–Gauss–Lobatto point distribution, and is allowed to exhibit discontinuities between elements in an L^2 -conforming sense. The spaces of shape functions are then given as

$$\mathcal{V}_h^u = \left\{ \mathbf{u}_h \in [L^2(\Omega_h)]^d : \mathbf{u}_h(\mathbf{x}^e(\boldsymbol{\xi}))|_{\Omega_e} = \tilde{\mathbf{u}}_h^e(\boldsymbol{\xi})|_{\tilde{\Omega}_e} \in \mathcal{V}_{h,e}^u = [\mathcal{Q}_k(\tilde{\Omega}_e)]^d, \forall e = 1, \dots, N_{\text{el}} \right\},$$

$$\mathcal{V}_h^p = \left\{ p_h \in L^2(\Omega_h) : p_h(\mathbf{x}^e(\boldsymbol{\xi}))|_{\Omega_e} = \tilde{p}_h^e(\boldsymbol{\xi})|_{\tilde{\Omega}_e} \in \mathcal{V}_{h,e}^p = \mathcal{Q}_{k-1}(\tilde{\Omega}_e), \forall e = 1, \dots, N_{\text{el}} \right\}.$$

Here, \mathcal{Q}_k denotes the space of tensor-product polynomials of degree k , and we highlight the ambiguity in notation related to the spatial wavenumber k in the context of energy spectra. The
 345 polynomial degree of the velocity shape functions is k , while it is $k - 1$ for the pressure for reasons of inf–sup stability. In this work, we only consider problems with Cartesian meshes so that the mapping $\mathbf{x}^e(\boldsymbol{\xi})$ is an affine transformation. At the interface between two elements e^- and e^+ , the numerical solution is not unique and we use the superscripts $-$ and $+$ to denote the solution from the two sides of an interface. When referring to element e , we denote interior information by $-$,
 350 and the outward pointing unit normal vector by $\mathbf{n} = \mathbf{n}^-$. To define numerical fluxes, we introduce the average operator $\{\{u\}\} = (u^- + u^+)/2$ and the jump operator $[[u]] = u^- \otimes \mathbf{n}^- + u^+ \otimes \mathbf{n}^+$, where $\mathbf{n}^+ = -\mathbf{n}^-$. We apply the usual abbreviations of integrals, $(v, u)_{\Omega_e} = \int_{\Omega_e} v \odot u \, d\Omega$ for volume integrals and $(v, u)_{\partial\Omega_e} = \int_{\partial\Omega_e} v \odot u \, d\Gamma$ for surface integrals with \odot denoting an inner product. We now state the weak formulation for all sub-steps of the projection scheme.

355 Obtain the numerical solutions $\hat{\mathbf{u}}_h, \hat{\hat{\mathbf{u}}}_h, \hat{\hat{\hat{\mathbf{u}}}}_h, \mathbf{u}_h^{n+1} \in \mathcal{V}_h^u$ and $p_h^{n+1} \in \mathcal{V}_h^p$ by testing with all test functions $\mathbf{v}_h \in \mathcal{V}_{h,e}^u$, $q_h \in \mathcal{V}_{h,e}^p$ for all elements $e = 1, \dots, N_{\text{el}}$:

In the first step, the convective term is discretised by the local Lax–Friedrichs flux

$$\left(\mathbf{v}_h, \frac{\gamma_0^n \hat{\mathbf{u}}_h - \sum_{i=0}^{J-1} \alpha_i^n \mathbf{u}_h^{n-i}}{\Delta t_n} \right)_{\Omega_e} =$$

$$- \sum_{i=0}^{J-1} \beta_i^n \left(- \left(\nabla \mathbf{v}_h, (\mathbf{u}_h \otimes \mathbf{u}_h)^{n-i} \right)_{\Omega_e} + \left(\mathbf{v}_h, \left(\{\{ \mathbf{u}_h \otimes \mathbf{u}_h \}\} + \frac{\Lambda}{2} [[\mathbf{u}_h]] \right)^{n-i} \cdot \mathbf{n} \right)_{\partial\Omega_e} \right), \quad (2.7)$$

where the stabilisation parameter is chosen as $\Lambda = \max(|\mathbf{u}_h^- \cdot \mathbf{n}|, |\mathbf{u}_h^+ \cdot \mathbf{n}|)$. The pressure Poisson operator is discretised by the symmetric interior penalty Galerkin (SIPG) method

$$\begin{aligned} & \left(\nabla q_h, \nabla p_h^{n+1} \right)_{\Omega_e} - \left(\nabla q_h, \frac{1}{2} \llbracket p_h^{n+1} \rrbracket \right)_{\partial\Omega_e} - \left(q_h, \{ \{ \nabla p_h^{n+1} \} \} \cdot \mathbf{n} \right)_{\partial\Omega_e} \\ & + \left(q_h, \tau \llbracket p_h^{n+1} \rrbracket \cdot \mathbf{n} \right)_{\partial\Omega_e} = - \frac{\gamma_0^n}{\Delta t_n} \left(- (\nabla q_h, \hat{\mathbf{u}}_h)_{\Omega_e} + (q_h, \{ \{ \hat{\mathbf{u}}_h \} \} \cdot \mathbf{n})_{\partial\Omega_e} \right), \end{aligned} \quad (2.8)$$

where a central flux is used for the velocity divergence operator on the right-hand side of the pressure Poisson equation. A central flux is also used for the pressure gradient term in the projection step

$$\left(\mathbf{v}_h, \hat{\mathbf{u}}_h \right)_{\Omega_e} = (\mathbf{v}_h, \hat{\mathbf{u}}_h)_{\Omega_e} - \frac{\Delta t_n}{\gamma_0^n} \left(- (\nabla \cdot \mathbf{v}_h, p_h^{n+1})_{\Omega_e} + (\mathbf{v}_h, \{ \{ p_h^{n+1} \} \} \mathbf{n})_{\partial\Omega_e} \right). \quad (2.9)$$

The viscous term is also discretised by the SIPG method

$$\begin{aligned} & \left(\mathbf{v}_h, \frac{\gamma_0^n}{\Delta t_n} \hat{\hat{\mathbf{u}}}_h \right)_{\Omega_e} + \left(\nabla \mathbf{v}_h, \nu \nabla \hat{\hat{\mathbf{u}}}_h \right)_{\Omega_e} - \left(\nabla \mathbf{v}_h, \frac{\nu}{2} \llbracket \hat{\hat{\mathbf{u}}}_h \rrbracket \right)_{\partial\Omega_e} \\ & - \left(\mathbf{v}_h, \nu \{ \{ \nabla \hat{\hat{\mathbf{u}}}_h \} \} \cdot \mathbf{n} \right)_{\partial\Omega_e} + \left(\mathbf{v}_h, \nu \tau \llbracket \hat{\hat{\mathbf{u}}}_h \rrbracket \cdot \mathbf{n} \right)_{\partial\Omega_e} = \left(\mathbf{v}_h, \frac{\gamma_0^n}{\Delta t_n} \hat{\hat{\mathbf{u}}}_h \right)_{\Omega_e}, \end{aligned} \quad (2.10)$$

while this step is skipped, $\hat{\hat{\mathbf{u}}}_h = \hat{\mathbf{u}}_h$, in the inviscid limit when solving the incompressible Euler equations. In a final postprocessing step, consistent divergence and continuity penalty terms are applied to weakly enforce the incompressibility constraint and normal continuity of the velocity field (Fehn *et al.* 2018b, 2021)

$$\begin{aligned} & \left(\mathbf{v}_h, \mathbf{u}_h^{n+1} \right)_{\Omega_e} + \left(\nabla \cdot \mathbf{v}_h, \tau_D \nabla \cdot \mathbf{u}_h^{n+1} \right)_{\Omega_e} \Delta t_n \\ & + \left(\mathbf{v}_h \cdot \mathbf{n}, \tau_C (\mathbf{u}_h^- - \mathbf{u}_h^+)^{n+1} \cdot \mathbf{n} \right)_{\partial\Omega_e} \Delta t_n = \left(\mathbf{v}_h, \hat{\hat{\mathbf{u}}}_h \right)_{\Omega_e}. \end{aligned} \quad (2.11)$$

The above postprocessing step is specifically related to the type of spatial discretisation used in this work, i.e., the L^2 -conforming discontinuous Galerkin method, and disappears in the continuous case. It was found that the divergence and normal-continuity penalty terms are crucial in terms of mass conservation and energy stability in order to obtain a method that is robust for large Reynolds numbers and coarse spatial resolutions, and we refer to Fehn *et al.* (2018b, 2019) for detailed numerical justification and validation of this approach. In this context, we note that alternatives to this stabilised approach exist, e.g., by using tailored finite element function spaces. For example, function spaces can be used that result in an $H(\text{div})$ -conforming (normal-continuous) velocity field and that is divergence-free in every point of the computational domain. The present L^2 -conforming approach does not fulfil these two properties exactly, but in a weak finite element sense. Definitions of the penalty parameters τ , τ_D , τ_C are given in Fehn *et al.* (2018b), where the default penalty factor $\zeta = 1$ is used in the present work unless specified otherwise. In terms of the occurrence of finite-time singularities and anomalous energy dissipation, the present work makes use of the argument that – due to the weighted residual formulation – the present discretisation can be applied to problems which lack regularity and for which the differential form of the equations is no longer an appropriate description. With respect to the implementation of the method, integrals in the weak form are evaluated by means of Gaussian quadrature, with consistent integration according to the 3/2-rule for the non-linear convective term (also denoted as polynomial dealiasing). Since we consider uniform Cartesian meshes (with elements of size h in all coordinate directions) in the present work, integrals are calculated exactly.

To obtain the size of the next time step within the adaptive time stepping scheme, we use the

12

following local CFL condition evaluated at all quadrature points q of element e

$$\Delta t = \min_{e=1,\dots,N_{el}} \left(\min_q \frac{\text{Cr}}{k^{1.5}} \frac{h}{\|\mathbf{u}_h\|} \Big|_{q,e} \right), \quad (2.12)$$

ensuring stability of the time integration scheme by selecting a Courant number $\text{Cr} < \text{Cr}_{\text{crit}}$. Note that – due to the CFL condition – an increase in spatial resolution by reducing h or increasing k automatically implies an increase in temporal resolution through a reduced time step size.

380 The present incompressible flow solver is implemented in C++ and forms a core module of the ExaDG (High-Order **D**iscontinuous **G**alerkin for the **E**xa-**S**cale) software, which is publicly available on [github](https://github.com).[†] The ExaDG software primarily makes use of the open-source finite element library deal.II (Arndt *et al.* 2021; Alzetta *et al.* 2018). The computational efficiency of the present discretisation methods in terms of fast implementations and fast iterative solvers is discussed in Fehn *et al.* (2018a, 2020); Kronbichler & Kormann (2019); Arndt *et al.* (2020).
385

3. Results

This section presents numerical results for three test cases in $d = 1, 2$, and 3 space dimensions. The one-dimensional problem is the well-known inviscid Burgers equation developing a singularity in finite-time for appropriate initial conditions. The two-dimensional example is a shear layer roll-up problem. The particular example used for the $d = 2$ investigations is not of primary importance as it is known theoretically that regularity is expected for two-dimensional Euler flows when starting from regular initial data. Instead, the aim of these one- and two-dimensional examples is an investigation to which extent the numerical discretisation scheme is able to mimic physical behaviour with potentially singular solutions. Having validated the numerical method for these well-understood problems, it is applied to the three-dimensional inviscid Taylor–Green problem, for which the physical understanding in terms of the occurrence of finite-time singularities and the related aspect of anomalous energy dissipation is speculative at present. As a further preparation, we summarise the quantities of interest in the following subsection.
395

3.1. Quantities of interest

400

Since turbulent flows in three space dimensions are our primary interest, we restrict the discussion in this subsection to the three-dimensional case implying extensions of certain relations to one- and two-dimensional problems only where possible. Of primary importance for the present study is the temporal evolution of the kinetic energy

$$E(t) = \frac{1}{V_\Omega} \int_\Omega \frac{1}{2} \mathbf{u}(\mathbf{x}, t) \cdot \mathbf{u}(\mathbf{x}, t) \, d\Omega,$$

and its dissipation rate dE/dt . The kinetic energy is normalised by the volume $V_\Omega = \int_\Omega 1 \, d\Omega$ of the computational domain. The integrals are evaluated numerically by means of Gaussian quadrature with $k + 1$ quadrature points in each coordinate direction. The time derivative used to obtain the dissipation rate is computed numerically from the kinetic energy at discrete instants of time via a second order finite difference formula for variable time step sizes with first order approximations at the end points. If anomalous dissipation ($dE/dt < 0$) occurs, the temporal evolution of the entropy \mathcal{E} ,

$$\mathcal{E}(t) = \frac{1}{V_\Omega} \int_\Omega \frac{1}{2} \boldsymbol{\omega}(\mathbf{x}, t) \cdot \boldsymbol{\omega}(\mathbf{x}, t) \, d\Omega,$$

[†] See <https://github.com/exadg/exadg>.

is expected to exhibit a singularity $\mathcal{E} \rightarrow \infty$ in finite time. Integrals are computed by Gaussian quadrature, which is exact down to round-off errors due to polynomial integrands and Cartesian meshes. A related local quantity is the maximum vorticity

$$\|\boldsymbol{\omega}\|_{\infty}(t),$$

where we take the maximum over all quadrature points over all elements in the discrete case and monitor its evolution over time with the interest in a detection of potentially singular behaviour in finite time. Although the maximum vorticity will remain finite for every numerical simulation of finite resolution, a mesh refinement study may give hints on the expected behaviour if the resolution was further increased. Finally, we consider kinetic energy spectra by transformation into wavenumber space \mathbf{k} (Cichowlas & Brachet 2005; Bustamante & Brachet 2012)

$$E(k, t) = \frac{1}{2} \lim_{\Delta k \rightarrow 0} \frac{\int_{\|\mathbf{k}\|=k}^{\|\mathbf{k}\|=k+\Delta k} \|\hat{\mathbf{u}}(\mathbf{k}, t)\|^2 d\mathbf{k}}{\Delta k} \stackrel{\text{DFT}}{\approx} \sum_{\substack{\mathbf{k} \in \mathbb{Z}^3 \\ k-\frac{1}{2} \leq \|\mathbf{k}\| < k+\frac{1}{2}}} \frac{1}{2} \|\hat{\mathbf{u}}_{\text{DFT}}(\mathbf{k}, t)\|^2,$$

where $\hat{\mathbf{u}}(\mathbf{k}, t)$ denotes the Fourier transform of the velocity, which only exists at discrete wavenumber vectors \mathbf{k} in case of a discrete Fourier transformation (DFT) obtained from sampled data of a discrete velocity field. The solution is first interpolated onto N equidistant points per element and per coordinate direction[†] to which the discrete Fourier transformation is applied, using the library FFTW (Frigo & Johnson 2005) in the present work. Note that considering $E(k, t)$ as a function of a scalar wavenumber k as well as the summation over spheres of radius k introduces the assumptions of homogeneity and isotropy. In case of anomalous energy dissipation, the enstrophy is expected to become infinite. Hence, exploiting the relation $\mathcal{E}(k, t) = k^2 E(k, t)$ between the enstrophy and energy spectra and further assuming a power law behaviour for the kinetic energy spectrum of the form $E(k, t) = C(t)k^{-n(t)}$, a singularity at time $t = t_*$ with $\mathcal{E}(t_*) = \int_0^{\infty} \mathcal{E}(k, t_*) dk = \infty$ would correspond to a decay with slope $n(t_*) = 3$ in the energy spectrum, see also Orlandi *et al.* (2012); Orlandi (2009). We use this criterion as a further validation of the results in case other quantities give hints of a potentially singular behaviour. Apart from that, energy spectra are typically investigated to assess the well-known $k^{-5/3}$ Kolmogorov spectrum for fully-developed, homogeneous isotropic turbulence. We apply this property to investigate whether the numerical results match expected physical behaviour obtained from classical cascade pictures in case of inviscid flows and beyond the time of potential singularities, $t > t_*$.

3.2. One-dimensional inviscid Burgers equations

We begin with studying the one-dimensional inviscid Burgers equation (Burgers 1948)

$$\frac{\partial u}{\partial t} + \frac{\partial}{\partial x} \frac{u^2}{2} = 0,$$

as a simplified model for the incompressible Euler equations. It is well known that this equation develops singularities in finite time, see for example Sulem *et al.* (1983); Dubrulle (2019). Previous works demonstrate that the use of energy-conserving discretisation schemes leads to thermalisation for this type of problems, see for example Ray *et al.* (2011); Murugan *et al.* (2020).

[†] The number of sampling points is chosen as $N = k + 1$ in the present work, i.e., equal to the number of nodal points of the discontinuous Galerkin discretisation, where k is the polynomial degree of the shape functions and should not be mixed up with the wavenumber k typically used in the context of energy spectra. The equidistant interpolation points all lie within the element away from the boundaries where the solution is discontinuous. If interpolation points on the boundary are used, one typically takes the average of the solution from neighbouring elements.

14




Figure 2: One-dimensional inviscid Burgers equations for two different initial solutions that form a singularity: On the left, the initial condition is a sine function, while it is a simple hat function that is piecewise linear on the right. The spatial resolution used for the computations corresponds to refinement level $l = 6$ and polynomial degree $k = 3$, resulting in an effective resolution of 256^1 .

It is therefore particularly interesting to study the behaviour of a discretisation scheme for this simplified problem first. The spatial discretisation is based on a discontinuous Galerkin scheme very similar to the one described in section 2 for the two- and three-dimensional case, i.e., the convective term is discretised with a local Lax–Friedrichs flux. Gaussian quadrature with a 3/2-overintegration rule is used as in the higher dimensional case due to the quadratic nonlinearity of the convective term, but intentionally no additional measure such as limiting, filtering, or other Riemann fluxes are taken to specifically address the jump that forms in the solution. We also emphasise that no artificial viscosity approach is used to deal with the singularity. For time integration, the classical explicit fourth-order Runge–Kutta method is used with a Courant number of $Cr = 0.4$.

Figure 2 shows the numerical solution $u_h(x)$ at various instants of time and the formation of a shock. The problem is solved on the domain $\Omega = [-1, 1]$ with Dirichlet boundary conditions prescribed at both boundary points of the one-dimensional domain. An equidistant grid with 2^l elements is used where l denotes the level of refinement. For polynomial approximations of degree k , the effective resolution becomes $(k + 1)2^l$. We exemplarily select two different initial solutions, a sine function, $u_h(x, t = 0) = -\sin(\pi x)$, and a hat function, $u_h(x, t = 0) = -2|x + 0.5| + 1$ for $x < 0$ and $u_h(x, t = 0) = 2|x - 0.5| - 1$ for $x \geq 0$. Due to the chosen initial conditions with $u > 0$ for $x < 0$ and vice versa, the solution piles up in the middle of the domain and a singularity ($\partial u / \partial x \rightarrow \infty$) forms at $x = 0$ in both cases. The oscillating behaviour of the numerical solution around the singularity could be improved by the advanced discretisation techniques mentioned above. From the results shown in figure 2 it is plausible that the kinetic energy is conserved until the formation of the shock and that energy will be dissipated at later times. For the hat function chosen as initial condition, it is straight-forward to derive an analytical expression for the temporal evolution of the kinetic energy as well as its dissipation rate, which is why we consider this setup in more detail in the following. According to the method of characteristics it follows that the shock forms at time $t_* = 0.5$. From that time on, the solution can be written as

$$u(x, t) = f(t) (x - \text{sign}(x)) ,$$

where $\text{sign}(x)$ takes values of ± 1 depending on the sign of the argument. The temporal evolution part $f(t)$ which describes the absolute value of u taken to the left and right of the origin of the




Figure 3: One-dimensional inviscid Burgers equations with hat function as initial condition: temporal evolution of kinetic energy as well as dissipation rate and convergence towards analytical profile for a mesh refinement study with refine levels $l = 2, \dots, 5$ and polynomial degree $k = 3$, resulting in effective resolutions of $16^1, \dots, 128^1$.

coordinate system at $x = 0$ can be obtained from the following consideration

$$f(t + dt) = f(t) - \underbrace{\frac{\partial u(x, t)}{\partial x} \Big|_{x=0^-}}_{=f(t)} \underbrace{dx}_{=u(x=0^-, t)dt} = f(t)(1 - f(t)dt),$$

i.e., the solution at $x = 0^-$ at time $t + dt$ equals the solution at position $-dx = -u(x = 0^-, t)dt$ at time t . Separation of variables and integration yields the result $f(t) = 1/(t + t_*)$. The kinetic energy $E(t) = \int_{\Omega} \frac{1}{2} u^2(x, t) dx$ is therefore given as

$$E(t) = \int_{-1}^1 \frac{1}{2} f^2(t) (x - \text{sign}(x))^2 dx = \frac{f^2(t)}{3}.$$

The kinetic energy dissipation rate is obtained by differentiation which yields a $(t + t_*)^{-3}$ decay for times $t \geq t_*$. The dissipation rate is $\Delta u^3/12$ when expressed in terms of the jump Δu of the solution, in agreement with the result in Dubrulle (2019) where it is noted that this inviscid dissipation is identical to the dissipation of the viscosity solution in the limit $\nu \rightarrow 0$. In figure 3, we show results for both the kinetic energy and the dissipation rate for a sequence of mesh refinement levels of $l = 2, \dots, 5$ with degree $k = 3$, resulting in effective resolutions of $16^1, \dots, 128^1$. For increasing spatial resolution, the numerical results converge to the analytical profiles. It can be seen that achieving grid-convergence for the dissipation rate requires higher spatial resolutions as compared to the temporal evolution of the kinetic energy itself. This is expected since the dissipation rate contains a temporal derivative that results in a higher sensitivity with respect to deviations (here numerical discretisation error) from the exact solution.

Figure 4 shows the same results for the problem with sine function as initial condition. We observe that the onset of energy dissipation is smooth as opposed to the hat function where the kinetic energy exhibits a kink and the dissipation rate a jump at the time of the singularity. In other words, the occurrence of a finite-time singularity does not imply an instantaneous onset of dissipation. We keep this in mind when considering the three-dimensional inviscid Taylor–Green problem which is a problem that also starts from sine-like initial data.

The point that we want to make with this example is that a discretisation scheme that involves purely numerical mechanisms of dissipation can provide the physically correct amount of dissipation for a sufficiently fine spatial resolution, see also the discussion in the introduction. Note that this is fundamentally different from viscosity solutions u^ν for small $\nu > 0$, for which the




Figure 4: One-dimensional inviscid Burgers equations with sine function as initial condition: temporal evolution of kinetic energy as well as dissipation rate and mesh refinement study with refine levels $l = 2, \dots, 5$ and polynomial degree $k = 3$, resulting in effective resolutions of $16^1, \dots, 128^1$.

455 required dissipation is realised by the additional viscous term in the equations and for which the
dissipation stemming from the numerical discretisation scheme tends to zero if the mesh resolves
the viscosity solution u^v exhibiting steep but finite gradients. Although the one-dimensional
Burgers equation can not reflect the complexity of three-dimensional turbulent flows, these results
put confidence in numerical discretisation schemes to also predict the solution in a physically
460 correct way for the higher-dimensional problems studied below.


3.3. Two-dimensional shear layer problem

We consider the two-dimensional shear layer roll-up problem (Brown 1995) where the initial velocity is given as


$$\mathbf{u}(\mathbf{x}, t = 0) = (\tanh(\rho(0.25 - |x_2 - 0.5|)), \delta \sin(2\pi x_1))^T.$$

Following Brown (1995), we set the two parameters ρ, δ to $\rho = 30$ and $\delta = 0.05$. The problem is solved on the domain $\Omega = [0, 1]^2$ with periodic boundaries in both directions. In the following, different viscous simulations with viscosities $\nu = 2.5 \cdot 10^{-3}, 10^{-3}, 10^{-4}$ are considered, as well as
465 the inviscid limit with $\nu = 0$. The mesh is uniform Cartesian with $(2^l)^2$ elements for refinement level l , the polynomial degree of the shape functions is $k = 7$, resulting in the effective resolution of $((k + 1)2^l)^2$. The simulations are run for the time interval $0 \leq t \leq 4$. The time step size is adapted dynamically with a Courant number of $\text{Cr} = 0.25$.

The aim of this example is to verify the robustness and accuracy of the present high-order
470 discontinuous Galerkin discretisation for a simple two-dimensional example. As mentioned in the introduction, the energy is conserved for the two-dimensional incompressible Euler equations, and this property should be preserved by a consistent discretisation scheme for sufficiently fine spatial resolutions. Figure 5 shows contour plots of velocity magnitude and vorticity magnitude at time $t = 1.2$ for a mesh with 16^2 elements (refinement level $l = 4$) for different values of the viscosity. In figure 6, we show the temporal evolution of the kinetic energy and the kinetic energy
475 dissipation rate for the different viscosity values. For each viscosity, results obtained on three meshes of increasing resolution with $4^2, 8^2$, and 16^2 elements are shown. For large viscosities, $\nu = 2.5 \cdot 10^{-3}$ and 10^{-3} , the results for the temporal evolution of the kinetic energy and dissipation rate coincide for all meshes. Also for the smallest viscosity of $\nu = 10^{-4}$ and the inviscid limit $\nu = 0$ the results obtained on the two finest meshes coincide and only minor
480 deviations can be observed for the coarsest mesh. This is in qualitative agreement with the contour



(a) Velocity magnitude for $\nu = 2.5 \cdot 10^{-3}, 10^{-3}, 10^{-4}, 0$ (from left to right).



(b) Vorticity magnitude for $\nu = 2.5 \cdot 10^{-3}, 10^{-3}, 10^{-4}, 0$ (from left to right).

Figure 5: Two-dimensional shear layer roll-up problem: contour plots of velocity magnitude and vorticity magnitude at time $t = 1.2$ for four different values of the viscosity (blue indicates low value and red high value). The results shown correspond to a mesh with 16^2 elements with a polynomial degree of $k = 7$ (effective resolution 128^2).

plots for the velocity magnitude in figure 5 which demonstrate that the velocity field is smooth and well-resolved on the finest mesh for all viscosities. The resolution requirements are higher for the vorticity containing spatial derivatives of the velocity field. As already noted in Brown (1995), the vorticity field is still not well-resolved even if convergence has already been achieved for the velocity or kinetic energy. It can be seen from figure 5 that the vorticity field is well-resolved for the viscous cases $\nu = 2.5 \cdot 10^{-3}, 10^{-3}, 10^{-4}$, but shows grid-dependence with numerical artefacts in the form of elevations of the vorticity at the element corners especially in the thin shear layer that is most difficult to resolve. In agreement with what is expected physically, the kinetic energy dissipation rate tends to zero for $\nu \rightarrow 0$ and the kinetic energy is conserved in the inviscid limit $\nu = 0$. Of particular importance w.r.t. the interpretation of results shown in section 3.4 for the three-dimensional Taylor–Green problem is the observation that the numerical dissipation occurring in the inviscid limit $\nu = 0$ for coarse spatial resolutions decreases to zero under mesh refinement for this two-dimensional problem.

An important aspect concerns the numerical robustness of the discretisation scheme. In Chalmers *et al.* (2019), instabilities are reported for the same shear layer problem with viscosity $\nu = 0$ for a discontinuous Galerkin discretisation with polynomial degree $k = 7$ and refinement level $l = 4$. This originates from the fact that the stabilised discretisation techniques developed in Krank *et al.* (2017); Fehn *et al.* (2018b) that render the discretisation robust in under-resolved scenarios and that are used in the present work have not been applied in that study. No robustness problems have been observed for the present discretisation scheme even for the coarsest resolutions, e.g. refinement levels of $l = 0, 1$ not shown explicitly here, which is a prerequisite to obtain a feasible incompressible flow solver for three-dimensional turbulent flow problems that are even more challenging in terms of the stability of a numerical discretisation scheme. Instabilities have also been reported for continuous spectral element discretisations for this two-dimensional shear layer problem, where filtering techniques can be used to recover stability (Fischer & Mullen 2001), at the cost of introducing new parameters into the discretisation scheme. A recent study by Thalabard *et al.* (2020) uses a pseudo-spectral

18




Figure 6: Two-dimensional shear layer roll-up problem: temporal evolution of kinetic energy and kinetic energy dissipation rate for decreasing viscosity values of $\nu = 2.5 \cdot 10^{-3}$, 10^{-3} , 10^{-4} , and 0. For each viscosity, results are shown for three different effective resolutions of 32^2 (dotted lines), 64^2 (dashed lines), and 128^2 (solid lines) corresponding to meshes with 4^2 , 8^2 , and 16^2 elements with polynomial degree $k = 7$.

510 scheme with hyperviscous linear dissipation to investigate a two-dimensional Kelvin–Helmholtz problem. A discretisation technique with properties similar to the present stabilised DG
 discretisation in terms of robustness and accuracy are exactly divergence-free $H(\text{div})$ -conforming
 discretisations, see for example the studies by Guzmán *et al.* (2016); Fu (2019) analysing this
 shear-layer problem, the study by Schroeder & Lube (2018) discussing other two-dimensional
 515 examples such as the Kelvin–Helmholtz instability problem, and the study by Fehn *et al.* (2019)
 comparing $H(\text{div})$ - and stabilized L^2 -conforming discretisations for three-dimensional turbulent
 flow problems in under-resolved scenarios.

3.4. Three-dimensional Taylor–Green vortex problem

We consider the 3D Taylor–Green vortex problem (Taylor & Green 1937) defined by the following initial velocity field

$$\mathbf{u}(\mathbf{x}, t = 0) = (\sin x_1 \cos x_2 \cos x_3, -\cos x_1 \sin x_2 \cos x_3, 0)^T.$$

Results of viscous simulations for increasing Reynolds number $\text{Re} = \frac{1}{\nu}$ are given in figure 1. In the following, the focus is entirely on the inviscid limit $\nu = 0$. The simulations are run over the time interval $0 \leq t \leq T = 20$ to cover the different flow regimes of laminar flow, transition to turbulence, and decaying turbulence. To reduce computational costs for fixed resolution of the flow (or to increase the effective resolution for a given amount of computational costs) it is common practise to exploit the symmetry of the Taylor–Green problem and simulate the flow on the impermeable box $\Omega = [0, \pi]^3$ with symmetry boundary conditions on all boundaries (Brachet *et al.* 1983), i.e.,

$$\mathbf{u} \cdot \mathbf{n} = 0, \quad \frac{\partial \mathbf{u}}{\partial n} = 0,$$

520 as opposed to the periodic box $\Omega = [-\pi, \pi]^3$ that is also used in computational studies. This optimisation allows to reduce computational costs by a factor of 8 for the present DG discretisation. Further symmetries can be exploited by spectral methods leading to the so-called fundamental box (Brachet *et al.* 1983; Kida 1985).


The computational domain $\Omega = [0, \pi]^3$ is discretised using a uniform Cartesian grid consisting of $(2^l)^d$ elements, where l denotes the level of refinement. The number of unknowns is given as $N_{\text{DoFs}} = (2^l)^d (d(k_u + 1)^d + (k_p + 1)^d) = (2^l)^d (d(k + 1)^d + k^d)$. It is common practise

525 in the literature to express the effective mesh resolution in terms of the periodic box to obtain comparability between different discretisation techniques that exploit different levels of symmetry. Hence, we define the effective spatial resolution as $(2^{l+1}(k+1))^d$, e.g., the effective resolution is 64^3 for refine level $l = 3$ and polynomial degree $k = 3$. Absolute tolerances of 10^{-12} and relative tolerances of 10^{-6} are used for the iterative linear solvers, where relative tolerance means
 530 that the residual is reduced by a factor of 10^{-6} compared to the initial residual that uses as initial guess a high-order extrapolation of the solution from previous time steps. The polynomial degree used for the TGV simulations is $k = 3$ and adaptive time stepping with $\text{Cr} = 0.25$ is used for all simulations. The penalty factors of the divergence and continuity penalty terms are chosen as defined in Fehn *et al.* (2018b), except for the finest resolution of 8192^3 where the penalty
 535 factors have been increased by a factor $\zeta = 2$ compared to the standard definition. For this fine resolution, the simulation remained stable also for the default value of $\zeta = 1$, but we observed oscillations in the maximum vorticity at early times, giving an indication of the need for a slightly larger penalisation of the divergence-free constraint and normal continuity of the velocity field. Since these oscillations disappeared when increasing the penalty factors by a factor of 2, this
 540 value has finally been used for this highest resolution simulation. The highest resolution of 8192^3 has $N_{\text{DoFs}} = 2.35 \cdot 10^{11}$ unknown degrees of freedom, and $2.27 \cdot 10^5$ time steps have been solved during this simulation. The computations have been performed on a large supercomputer using almost 100k cores for the highest resolution, requiring a run time of approximately 8.4 days.


3.4.1. Recapitulating the state-of-the-art

545 This section briefly summarises the type of discretisation, the maximal effective resolutions, and the final time T of the simulations considered in previous numerical studies for the three-dimensional inviscid Taylor–Green vortex problem. In Brachet *et al.* (1983), a spectral method with maximum resolution of 256^3 (exploiting symmetry) has been used and direct simulation has been performed up to times $t \leq 4$. In a subsequent work by Brachet *et al.* (1992), a maximum
 550 resolution of 864^3 (exploiting symmetry) has been reached, and again direct simulation has been performed up to times $t \leq 4$. A comparison of a spectral method and WENO finite difference method can be found in Shu *et al.* (2005), where a maximum resolution of 368^3 (exploiting symmetry) and simulation up to times $t \leq 6$ has been considered. A modal discontinuous Galerkin method has been studied in Chapelier *et al.* (2012), with the simulations performed
 555 up to times $t \approx 5 - 7$ until the simulations became unstable, with the maximum resolution of around 96^3 for different polynomial degrees from $k = 1$ to $k = 5$. DG discretisations used to study the inviscid TGV problem have also been analyzed in Moura *et al.* (2017b,a); Fernandez *et al.* (2018); Manzanero *et al.* (2020); Schroeder (2019), but with a focus on LES (large eddy simulation) modelling. The study by Cichowlas & Brachet (2005) used a spectral method with
 560 maximum resolution of 2048^3 with simulations performed up to times $t \leq 4$. The highest resolution of 4096^3 has been achieved in Bustamante & Brachet (2012) using a spectral method, and the simulations have been performed up to times $t \leq 4$.

In these works, indications of finite-time singularities are mentioned. In Morf *et al.* (1980), $t_* = 5.2$ is obtained from power series expansions. A more accurate variant using power series
 565 expansions presented in Brachet *et al.* (1983) leads to $t_* = 4.4 \pm 0.2$. Furthermore, the study by Brachet *et al.* (1983) reports indirect evidence for a finite-time singularity according to the direct numerical simulation results but the authors conclude that the resolution of 256^3 is not sufficient to investigate the occurrence of finite-time singularities for times $t \geq 4$. The more recent study by Bustamante & Brachet (2012) estimates a blow-up time of $t_* \approx 4$ and concludes that the
 570 results are not inconsistent with the occurrence of a singularity. The work by Larios *et al.* (2018) obtains a blow-up time of $t_* \approx 4.2$ similar to the blow-up time in Brachet *et al.* (1983).



(a) Velocity magnitude on plane $x = \pi$ at times $t = 0, 1, 2, 3, 4$ (from left to right).



(b) Vorticity magnitude on plane $x = \pi$ at times $t = 0, 1, 2, 3, 4$ (from left to right).

Figure 7: Three-dimensional inviscid Taylor–Green problem: contour plots of velocity magnitude and vorticity magnitude (blue indicates low value and red high value) on plane $x = \pi$ of impermeable box. The results shown correspond to a mesh with 32^3 elements with a polynomial degree of the shape functions of $k = 3$ (effective resolution 256^3).

3.4.2. Results in physical space

The early stage of the Taylor–Green vortex evolution with the formation of thin flow structures is visualised in figure 7. Similar results have been shown and discussed in detail already in Brachet *et al.* (1983) for the same effective resolution of 256^3 using a spectral method. In agreement with the above results for the two-dimensional shear layer problem, we observe that the velocity field is resolved at all times, while under-resolution effects are clearly visible in the contour plots of the vorticity magnitude at later times $t = 3$ and $t = 4$ for the chosen resolution. A high-resolution visualisation of the thin vortex sheet shown in figure 7 with a volume rendering of the vorticity magnitude has been shown in Bustamante & Brachet (2012) for an effective resolution of 4096^3 . Figure 8 shows visualisation results for a high-resolution simulation (effective resolution 2048^3) at later times $t = 4, 5, 6, 7, 8, 9$, around which small-scale features occur and transition to turbulence takes place. These results illustrate that the present discretisation scheme does not lead to thermalisation, which appears to be a prerequisite to obtain physically meaningful results, compare for example the present results to the thermalised results shown in (Schroeder 2019, Figure 9.14). Flow visualisation is discussed in the literature as one possibility to trace finite-time singularities, but appears to be impractical due to the difficulties in handling large data sets for high-resolution simulations necessary for such investigations, and due to the difficulties in visualising singularities (that do not show up as singularities for a finite-resolution numerical simulation). Hence, the attention is turned to other techniques in the following.

We present numerical results of a mesh convergence study for refinement levels $l = 3, \dots, 10$ and polynomial degree $k = 3$. Figure 9 shows the temporal evolution of the kinetic energy and the kinetic energy dissipation rate. At small times t , the energy is constant and the energy dissipation rate is zero. This agrees with the expected theoretical behaviour stating energy conservation as long as the solution remains smooth and has also been shown in previous works in a similar way. In this work, we do not want to terminate the simulations once we expect the simulation to become under-resolved, but instead continue the simulations until $t = 20$. Depending on the effective mesh resolution, an onset of energy dissipation can be observed that shifts towards later




Figure 8: Three-dimensional inviscid Taylor–Green problem: iso-surfaces of Q-criterion at times $t = 4, 5, 6, 7, 8, 9$, where the blue surface corresponds to a value of -0.5 and the orange surface to a value of 0.5 . The results correspond to a mesh with 256^3 elements with a polynomial degree of the shape functions of $k = 3$ (effective resolution 2048^3).

times for increasing spatial resolution. However, this time of onset of dissipation does not seem to be pushed beyond $t \approx 5$ even for the finest spatial resolutions. This is illustrated more clearly in figure 10, which plots the kinetic energy dissipation rate in logarithmic scaling as well as the error against the fine-resolution simulation (ref). The fact that the dissipation rate of the simulations with 2048^3 , 4096^3 resolution “converges” to that of the finest resolution at a time $t \approx 5$ for 2048^3 and $t < 5$ for 4096^3 is consistent with a potential blow-up time $t_* < 5$ in our opinion. Overall, we make the interesting phenomenological observation that the kinetic energy evolution and its dissipation rate tend to converge to a dissipative solution rather than an energy-conserving solution with vanishing dissipation rate. As in high-Reynolds number viscous simulations of this problem, the kinetic energy dissipation rate reaches a maximum at $t = 8 - 9$ and decreases afterwards. In section 3.4.5, grid-convergence of the sequence of discrete solutions to a dissipative reference solution is investigated in more detail. Note that the present results for the dissipation rate agree well with those shown in Cichowlas *et al.* (2005) for an energy-conserving scheme, where an effective dissipation rate is deduced from the thermalised energy $E_{\text{th}}(t)$, the energy associated to the small scales with wavenumber $k > k_{\text{th}}$, where k_{th} is the wavenumber at which the energy spectrum exhibits a local minimum.

It is now examined whether this behaviour is consistent with the temporal evolution of the maximum vorticity and the enstrophy shown in figure 11. For both quantities, one can immediately identify three phases: (i) a first phase up to approximately $t \approx 3$ in which the flow is well resolved for all spatial resolutions so that the results essentially overlap for all simulations, (ii) an intermediate phase $3 < t < 5$ in which the different simulations start to deviate from each other due to under-resolution effects depending on the spatial resolution of each simulation,

22




Figure 9: Three-dimensional inviscid Taylor–Green problem: temporal evolution of kinetic energy (left) and kinetic energy dissipation rate (right) for increasing effective spatial resolution.




Figure 10: Three-dimensional inviscid Taylor–Green problem: temporal evolution of kinetic energy dissipation rate in logarithmic scaling (left) and error against fine-resolution simulation (right).

and (iii) a final phase $t > 5$ in which the results of all simulations deviate substantially due to the different resolution capabilities of the different simulations. In the first phase, the vorticity first decreases, reaches a minimum, and then begins to grow exponentially in agreement with the results shown in Bustamante & Brachet (2012, figure 1 (b)). In figure 11, we added a reference curve with $\exp\left(\frac{2}{3}t\right)$ growth which describes the growth of the maximum vorticity very well in this regime. In the second phase at around $t \approx 3.5$, the maximum vorticity begins to grow substantially faster, and the growth of vorticity essentially depends on the spatial resolution that is directly linked to the maximum velocity gradient that can be represented on a given mesh. As already mentioned in the introduction, for every simulation that does not blow up due to numerical instabilities of a discretisation scheme, the maximum vorticity will remain finite no matter how fine the spatial resolution is. Therefore, the occurrence of a finite-time singularity with $\lim_{t \rightarrow t_*} \|\omega\|_\infty = \infty$ remains speculative. While our results might be considered consistent with such a vorticity blow-up scenario, we are not able to identify a concrete blow-up time $t = t_*$ from the present results. We rather observe that the time of maximum vorticity observed in this second phase is shifted to later times also for the finest spatial resolutions. At the same time, one might argue that a finite-time blow-up at a time $t = t_*$ with $t_* \approx 4 - 5$ would produce results similar to those shown here, with the maximum vorticity following the exact profile until the




Figure 11: Three-dimensional inviscid Taylor–Green problem: temporal evolution of maximum vorticity (left) and enstrophy (right) for increasing effective spatial resolution.

curve of a specific spatial resolution branches off due to under-resolution of the simulation. In such a scenario, one would expect the maximum vorticity to grow by a factor of 2 for refinement level $l \rightarrow l + 1$ due to the mesh size being reduced by a factor of 2, allowing numerical gradients becoming twice as large. While we observe such an increase in maximum vorticity from one refinement level to the next, it is not clear whether this suspected blow-up would happen in finite time. In the third phase, the maximum vorticity reaches a global maximum between $t = 6 - 7$ for each resolution before it starts to decrease slowly. In this phase, the maximum vorticity is offset by a factor of approximately 2 from one refine level to the next. This is a clear indication that none of the simulations is able to resolve the finest structures, and it is plausible that a factor of 2 in mesh size also gives a factor of 2 in maximum vorticity. Finally, the maximum vorticity shows a strongly fluctuating behaviour in the third phase. Note that the maximum vorticity is determined numerically by taking the maximum over all quadrature points, i.e., the vorticity field is sampled in discrete points. This effect is negligible for well-resolved scenarios but might explain an oscillatory behaviour in case a local maximum travels through the domain.

The temporal evolution of the enstrophy is overall similar to that of the maximum vorticity. An important difference is that the enstrophy does not reach a local minimum at early times as observed for the maximum vorticity. In the second phase, the growth of enstrophy is more moderate compared to the maximum vorticity. In the third phase, the enstrophy curves can be described essentially as smoothed variants of the maximum vorticity from which the high-frequency content has been removed. A possible explanation for the enstrophy behaviour in the second and third phase is that the enstrophy is not a local quantity, but an average in space over the computational domain. Again, a growth in enstrophy by a factor of two from one mesh level to the next is observed at later times, which is consistent with an enstrophy evolution theoretically taking infinite values, or taking finite values but much larger than those obtained numerically in figure 11. Considering Onsager’s conjecture as valid, it is clear that one can not expect convergence for the temporal evolution of maximum vorticity and enstrophy. Theoretically, convergence can then only be expected for the kinetic energy evolution and to some extent for kinetic energy spectra up to the resolution limits of the discretisation scheme, as discussed in the following.

3.4.3. Results in spectral space

Figure 12 shows kinetic energy spectra for increasing spatial resolution at various instants of time, namely at $t = 1, \dots, 9$ in steps of width 1. Results are shown for resolutions of 128^3 to 2048^3 . High computational costs and memory requirements of the FFT (fast fourier transformation) part of our simulations did not allow us to perform the spectral analysis for the highest resolutions




(a) Effective resolution of 128^3 (b) Effective resolution of 256^3 (c) Effective resolution of 512^3

Figure 12: Three-dimensional inviscid Taylor–Green problem: kinetic energy spectra for effective resolutions of 128^3 , 256^3 , 512^3 at times $t = 1, 2, \dots, 9$.

of 4096^3 and 8192^3 . For a discussion of the general behaviour of energy spectra as a function of time regarding the early time behaviour $t \leq 4$ we refer to Brachet *et al.* (1983); Cichowlas & Brachet (2005); Bustamante & Brachet (2012), where it is shown how the energy spectra can be fitted to functions of the form $E(k, t) = C(t)k^{-n(t)} \exp(-2k\delta(t))$ and where values obtained for $n(t)$ and $\delta(t)$ are discussed in detail. To verify the present results, we include reference



(d) Effective resolution of 1024^3 (e) Effective resolution of 2048^3

Figure 12: Three-dimensional inviscid Taylor–Green problem: kinetic energy spectra for effective resolutions of 1024^3 , 2048^3 at times $t = 1, 2, \dots, 9$.

law) or $n = 7/3$ (motivated by results obtained in Piatkowski (2019) for viscous Taylor–Green simulations). The energy spectra are compared against the slope $n = 3$ as a means to investigate the plausibility of a potentially singular behaviour and to identify a time $t = t_*$ at which such a blow-up could occur, as motivated in section 3.1. Once the flow has transitioned to a fully turbulent state, it can be expected that the energy spectrum exhibits some form of Kolmogorov scaling. For this purpose, we consider the energy spectra at times $t = 8$ and 9 where the maximum dissipation rate occurs, even though the flow might not be fully homogeneous isotropic at that time and the results might deviate from Kolmogorov’s $k^{-5/3}$ scaling. To quantify the resolution capabilities of the present discretisation, we also plot the Nyquist wavenumber k_{Nyquist} as well as the wavenumber $k_{1\%}$ according to the 1%-rule by Moura *et al.* (2017b) that aims at obtaining an accurate resolution limit for upwind-like DG discretisations for a specific polynomial degree of the function space.

Figure 12 shows that the range of scales resolved by the numerical method increases with increasing spatial resolution as expected theoretically, and that the resolution limit for polynomial degree 3 is described very well by the 1%-rule corresponding to this degree. The energy spectra reach a slope of -3 between $t = 4$ and $t = 5$, making our results consistent with the occurrence of singular behaviour around that time ($4 < t_* < 5$), in agreement with the rapid growth of the maximum vorticity observed in the same time interval, see figure 11. The spectra in the inertial range show a decay slightly stronger than $k^{-5/3}$ and better agreement is achieved when compared to a $k^{-7/3}$ scaling that is also shown in figure 12. This behavior has already been observed in Piatkowski (2019) for viscous Taylor–Green vortex simulations, where it was found

that Kolmogorov's $k^{-5/3}$ scaling can only be observed at later times, e.g., $t \approx 20$. Regarding the inertial scaling, the present results are therefore in agreement with results shown in the literature. Towards the Nyquist wavenumber, a moderate pile-up of energy can be observed by comparison against the $-5/3$ and $-7/3$ reference slopes before the energy falls off very rapidly. The energy pile up is characteristic of this type of high-order discontinuous Galerkin approach and becomes stronger for higher polynomial degrees, see Moura *et al.* (2017*b,a*) and references therein. This particular behaviour is often the primary target when optimizing discretisation schemes, see for example the recent studies by Flad & Gassner (2017); Winters *et al.* (2018); Manzanero *et al.* (2020) and references therein, where it is suggested to counteract this energy bump behaviour by explicit sub-grid scale modelling. However, the recent study by Fernandez *et al.* (2018) shows that this topic is delicate and improvements in some quantities by the use of explicit sub-grid models cause deviations in other quantities such as the temporal evolution of the kinetic energy dissipation rate. In our opinion, the challenge lies in improving the spectral behaviour and at the same time not giving up the improved resolution capabilities (per degrees of freedom) of high-order discretisations. The overall goal can be formulated as achieving a discretisation method that is accurate w.r.t. both the spectral behaviour and the behaviour in physical space, e.g., the temporal evolution of the kinetic energy and its dissipation rate. We want to note that from such a holistic view it is not clear a priori whether an explicit sub-grid scale model optimizing the energy spectrum according to the inertial $k^{-5/3}$ law is advantageous overall. We take up this point again in the outlook in section 4 discussing possible directions of future research. The energy spectra shown in (Schroeder 2019, figure 9.15) for an exactly energy-conserving discretisation scheme illustrate that such a scheme leads to physical inconsistencies for the $E(k)$ -curves in the inertial range.

3.4.4. Does the numerical dissipation have artificial or predictive character?

The dissipation of kinetic energy observed for the inviscid Taylor–Green simulations originates from the numerical method. At first sight, one might argue that changing the discretisation scheme by choosing another numerical flux or varying certain parameters leads to results that are more or less dissipative, i.e., that the amount of dissipation is artificial and is determined by the discretisation parameters. The results of the mesh convergence study shown above do not support this point of view, and we want to provide further results that might allow insights into the predictive character of these dissipative numerical solutions. In this context, it is illustrative to study the temporal evolution of the kinetic energy and its dissipation rate under a variation of parameters of the discretisation scheme. Figure 13 shows a parameters study of the penalty factor ζ of the divergence and continuity penalty terms of the present discretisation, considering values of $\zeta = 1, 2, 4, 8$ by the example of the 1024^3 spatial resolution. Note that this parameter is the crucial one in stabilizing the method in the under-resolved and high-Reynolds regime (Fehn *et al.* 2018*b*, 2019). We observe that the overall amount of dissipation as well as the dissipation maximum are essentially unaltered by a variation of this parameter. It is worth noting that the time of onset of dissipation is also not affected by the value of ζ . This might have important implications. Assuming that a non-dissipative (energy-conserving) solution is the correct physical behaviour and that the numerical dissipation is artificial, one might expect that an increase of the penalty parameter affects the numerical solution more strongly, e.g., changes the amount of overall dissipation or leads to a delayed onset of dissipation due to a better fulfilment of the divergence-free constraint for example. As we do not observe such a behaviour, these results are an indication that the obtained dissipative solutions have a predictive character in our opinion. It is clear that we can not expect the results to be identical for different penalty factors, since the results are not grid-converged for the chosen 1024^3 resolution, meaning that the discretisation scheme and its parameters affect the numerical solution.




Figure 13: Three-dimensional inviscid Taylor–Green problem: variation of penalty factor ζ and its influence on the temporal evolution of the kinetic energy (left) and the kinetic energy dissipation rate (right) for effective resolution of 1024^3 .

3.4.5. Are the results grid-converged?

We finally address the question whether and to which extent the results presented for inviscid Taylor–Green simulations can be considered as grid-converged. For this purpose, we compute relative L^2 -errors for the kinetic energy evolution and the kinetic energy dissipation rate

$$e_E^2 = \frac{\int_{t=0}^T (E(t) - E_{\text{ref}}(t))^2 dt}{\int_{t=0}^T (E_{\text{ref}}(t))^2 dt}, \quad e_{dE/dt}^2 = \frac{\int_{t=0}^T \left(\frac{dE(t)}{dt} - \frac{dE_{\text{ref}}(t)}{dt} \right)^2 dt}{\int_{t=0}^T \left(\frac{dE_{\text{ref}}(t)}{dt} \right)^2 dt}.$$

Since no analytical solution is available, the errors are measured using the finest resolution of 8192^3 as a reference (ref). This implies that we can not compute the error for the 8192^3 resolution. The error of this simulation can only be roughly estimated by extrapolating the convergence trend observed for the coarser resolutions and assuming that this convergence behaviour continues for the finest resolution. Defining a simulation with a relative error of 1% or less as grid-converged, the results in figure 14 reveal that we achieve grid-convergence in the kinetic energy evolution with errors below 1%. For the second finest resolution (the last data point in figure 14), the measured error is 0.27%. For the kinetic energy dissipation rate, the error is significantly larger, demonstrating that this quantity is more sensitive, in agreement with what has been observed in figure 1. For the second finest resolution, the measured error is 3.52%. While the errors can be expected to be smaller for the finest resolution 8192^3 that is used as a reference solution here, we conclude that even finer resolutions would be required to achieve grid-convergence also for the kinetic energy dissipation rate in terms of the 1% error level. Figure 14 also shows linear least-squares fits (e.g., $\log(E) \approx a \log(N_{\text{DoFs}}) + b$ for the kinetic energy) to the data obtained from the numerical experiments, where a mean converge rate of approximately $h^{3/4}$ is obtained for the kinetic energy and $h^{1/2}$ for the dissipation rate. This result – seemingly providing numerical evidence of grid-convergence to a dissipative solution of the three-dimensional incompressible Euler equations under Taylor–Green initial conditions – is the main result of the present study. While one might conjecture that this solution could be a weak Euler solution, there is currently no rigorous mathematical theory guaranteeing convergence to a weak solution. Moreover, even if we in fact obtained convergence to a weak Euler solution, it might not be the viscosity solution for $\nu \rightarrow 0$. Finally, from the results in figure 14 it can not be excluded that the dissipation rate would (slowly) tend to zero in the limit $h \rightarrow 0$ if finer spatial resolutions were realised. Let us explain this decisive aspect in more detail. For a solution that is non-dissipative (i.e., energy-conserving) from a physical perspective, one might argue that dissipation




Figure 14: Three-dimensional inviscid Taylor–Green problem: relative L^2 -errors of the temporal evolution of the kinetic energy (left) and the kinetic energy dissipation rate (right) for effective resolutions ranging from 8^3 to 4096^3 .

in the numerical simulation can only originate from under-resolution and can be expected to tend to zero when increasing the resolution. However, a dissipation rate decreasing extremely slowly under mesh refinement, e.g. as $1/\log \log(1/h)$, might then (erroneously) indicate grid-convergence to a dissipative solution, although the true solution is actually energy-conserving. Against this background, our numerical investigations according to the indirect approach are consistent with or suggestive of finite-time singularities, but do not conclusively provide evidence of finite-time singularities.

4. Conclusion and outlook

Hunting for finite-time singularities in incompressible Euler flows is a challenging discipline. Searching for singular behaviour in visualisations of three-dimensional simulation results evokes the picture of finding a needle in a haystack, as it can be expected that singularities are very localised, will never be resolved by a numerical scheme, and the amount of data for large-scale three-dimensional simulations soon becomes cumbersome. For this reason, most previous studies focused on quantities such as the maximum vorticity, the analyticity strip width, fitting energy spectra to power law behaviour, and related blow-up criteria. The present work focuses on global quantities such as the temporal evolution of the kinetic energy, avoiding geometrical complexities in visualisation and striving for a clearer indication of singular behaviour given that it might be computationally less demanding to resolve the kinetic energy than the vorticity in numerical simulations. We call this approach *indirect* since it exploits the connection between singular behaviour and anomalous energy dissipation according to Onsager’s conjecture. A decisive point is that this technique requires suitable discretisation schemes that remain robust in the presence of singularities and provide mechanisms of dissipation in case no viscous dissipation is present, which is a challenge in itself. Then, the idea is that observing an energy dissipating behaviour for a sequence of mesh refinement levels provides insight into the physical dissipation behaviour of the problem under investigation.

We applied this technique to one-, two-, and three-dimensional problems. Results consistent with theory have been obtained in one and two space dimensions. Subsequently, this technique has been used to study the complex three-dimensional inviscid Taylor–Green problem, for which an energy dissipating behaviour consistent with the high Reynolds number limit of viscous simulations available in the literature has been observed. The present study measures grid-convergence to a dissipative, fine-resolution numerical solution for the three-dimensional

inviscid Taylor–Green problem with a measured relative L^2 -error of 0.27% for the kinetic energy and 3.52% for the kinetic energy dissipation rate. The results for the temporal evolution of the kinetic energy might therefore be considered as grid-converged and serve as a reference solution for future studies. Regarding the temporal evolution of the maximum vorticity and the enstrophy, we are able to resolve an increase of almost four orders of magnitude for both quantities. To the best of our knowledge, these are the highest resolution results published to date for the three-dimensional inviscid Taylor–Green vortex problem. Confidence is put into the reliability of the numerical results for this challenging three-dimensional problem by the circumstance that the numerical method applied here is a robust discretisation scheme that remains numerically stable in the inviscid limit for all spatial resolutions that have been investigated. This is an important prerequisite to draw conclusions about a potential physical blow-up. In contrast, a numerical simulation that blows up in finite time does not allow any conclusions, since the observed blow-up is due to numerical instabilities. In other words, a physical blow-up does not imply a numerical blow-up for a finite-resolution numerical simulation.

In summary, this work wants to raise the questions (i) to which extent these results are related to weak dissipative solutions of the incompressible Euler equations, (ii) to which extent these results can then be interpreted as a numerical confirmation of the energy dissipation anomaly, and (iii) whether these results would imply finite-time singularities according to Onsager’s conjecture. Let us emphasise that the present results do not formally prove convergence of the discrete velocity fields \mathbf{u}_h to a weak Euler solution \mathbf{u} . Even if convergence to a weak Euler solution was obtained for a subsequence h_j , another subsequence h_k might converge to a different weak Euler solution. Moreover, it can not be excluded that the observed “dissipation anomaly” is of numerical origin in the sense that the spatial resolutions considered here might be too coarse to adequately resolve an energy-conserving Euler solution. Nevertheless, the present study might complement theoretical and experimental works on anomalous energy dissipation and the related implications on finite-time singularities for three-dimensional incompressible Euler flows according to Onsager’s conjecture. As part of future work, the so-called $4/5^{\text{th}}$ -law (Duchon & Robert 2000) should be investigated numerically in order to substantiate the hypothesis of having found a dissipative anomaly. Finally, the present work would gain further theoretical support by a proof of convergence to generalised weak Euler solutions for DG discretisation schemes of the Euler equations in the limit $h \rightarrow 0$ (along subsequences).

Given the challenges of the proposed indirect approach, it appears to be natural to raise the question whether this technique has advantages over well-known direct techniques. Let us share some ideas why the indirect approach might be an attractive technique for the exploration of finite-time singularities. A key motivation for the indirect approach is that it might not be necessary to resolve the smallest scales of the flow in order to resolve the temporal evolution of the kinetic energy. This is based on the observation that resolving the vorticity field (direct approach) typically requires significantly finer resolutions than resolving the kinetic energy (indirect approach). When studying singularities by the direct approach, one needs to realise that resolving the smallest scales is not possible for finite spatial resolutions. The vorticity will take finite \dagger values at all times for a stable discretisation scheme, so that results of a single simulation do not serve as numerical evidence of a finite-time singularity. A key element of the indirect approach is a mesh refinement study where the quantities of interest (kinetic energy and its dissipation rate) do not blow up. In our opinion, this circumstance supports cross-solver validation very naturally. Assume for example that different spatial discretisation schemes would converge to the same dissipative, numerical solution (measured by some error norms as done in the present work). A cross-solver validation of the kinetic energy evolution or dissipation rate (indirect approach) might give confidence that the

\dagger In this context, the term *finite* means that the quantity of interest takes values much smaller than the “infinite” value defined by the maximum floating point number that can be represented on a computer.

850 obtained dissipative behaviour is not artificial (why should entirely different PDE solvers produce
“the same” artificial result with artificial dissipation?). On the contrary, it would be difficult for
each of these simulations to demonstrate a blow-up of vorticity in order to provide evidence of
finite-time singularities by direct techniques. All these solvers might show an increase in vorticity
by a large factor, but it is still not clear whether a finite-time singularity $\|\omega\|_\infty \rightarrow \infty$ occurs.
855 Of course, there is currently not sufficient data available in the literature to substantiate this
assumption regarding a cross-solver validation of our results. One aspect that we want to initiate
with the present contribution is exactly such a cross-solver validation based on space-averaged
results or other suitable statistical results. In summary, the argument in favour of the indirect
approach would then be a clearer *indication* of finite-time singularities due to relaxed resolution
860 requirements as compared to the direct approach.

Applicability of high-order DG discretisations to large-scale problems in turbulence research,
for which spectral methods are currently the state-of-the-art due to their accuracy and
computational efficiency, has been demonstrated. Finally, the present study gives indications
in terms of what a promising large-eddy simulation strategy might be, and contributes to the
865 long-lasting and difficult discussion on explicit versus implicit sub-grid scale models. High-order
discretisations that can be described as implicit LES have shown very promising results for
moderate Reynolds number flows, but it is often argued that such techniques can be expected
to finally need explicit sub-grid scale modelling once they are applied in the limit $Re \rightarrow \infty$.
The present work contributes to this discussion by investigating a high-order discontinuous
870 Galerkin discretisation without explicit model in the inviscid limit. Assuming that such an
implicit approach gives physical results in the inviscid limit, e.g., consistent with Onsager’s
conjecture on anomalous energy dissipation, it makes one more confident that such a method
is able to naturally account for more complex physical mechanisms in turbulence beyond K41
theory (Dubrulle 2019). Taking as an alternative an energy-conserving numerical method with
875 explicit sub-grid scale model, the anomalous energy dissipation has to be realised by the sub-grid
model. We therefore believe it is an interesting future research direction to take the results shown
in the present study for the inviscid Taylor–Green problem as a reference for further validation
or to perform comparative studies between explicit and implicit LES techniques for the highest
Reynolds number case, the inviscid limit.

880 **Supplementary material.** Reference data for the inviscid Taylor–Green vortex simulations
for resolutions of 2048^3 , 4096^3 , 8192^3 (kinetic energy) and 2048^3 (kinetic energy spectra) is
available as supplementary material.

885 **Acknowledgements.** The authors thank Gert Lube and Oliver Neumann for inspiring discussions
on this topic. The authors also thank Marc Brachet and Samridhhi Sankar Ray for helpful
comments on a preprint of this work. We further appreciate contributions by Nisarg Patel
who provided the high-resolution visualization results for the inviscid Taylor–Green problem,
and support by the staff of the Leibniz Supercomputing Centre. Finally, contributions by the
890 developer communities of the ExaDG and deal.II software projects are gratefully acknowledged.

Funding. The research presented in this paper was partly funded by the German Research
Foundation (DFG) under the project “High-order discontinuous Galerkin for the EXA-scale”
(ExaDG) within the priority program “Software for Exascale Computing” (SPPEXA), grant
895 agreement no. KR4661/2-1 and WA1521/18-1. The authors gratefully acknowledge the Gauss
Centre for Supercomputing e.V. (www.gauss-centre.eu) for funding this project by providing
computing time on the GCS Supercomputer SuperMUC-NG at Leibniz Supercomputing Centre
(LRZ, www.lrz.de) through project id pr83te.

900 **Declaration of Interests.** The authors report no conflict of interest.

Author ORCID. N. Fehn, <https://orcid.org/0000-0001-7693-692X>; M. Kronbichler, <https://orcid.org/0000-0001-8406-835X>; W. A. Wall, <https://orcid.org/0000-0001-7419-3384>

REFERENCES

- 905 AGAFONTSEV, D. S., KUZNETSOV, E. A. & MAILYBAEV, A. A. 2015 Development of high vorticity structures in incompressible 3D Euler equations. *Physics of Fluids* **27** (8), 085102.
- AINSWORTH, M. 2004 Dispersive and dissipative behaviour of high order discontinuous Galerkin finite element methods. *Journal of Computational Physics* **198** (1), 106 – 130.
- 910 ALZETTA, G., ARNDT, D., BANGERTH, W., BODDU, V., BRANDS, B., DAVYDOV, D., GASSMOELLER, R., HEISTER, T., HELTAI, L., KORMANN, K., KRONBICHLER, M., MAIER, M., PELTERET, J.-P., TURCK SIN, B. & WELLS, D. 2018 The deal.II library, version 9.0. *J. Numer. Math.* **26** (4), 173–184.
- ARNDT, D., BANGERTH, W., DAVYDOV, D., HEISTER, T., HELTAI, L., KRONBICHLER, M., MAIER, M., PELTERET, J.-P., TURCK SIN, B. & WELLS, D. 2021 The deal.II finite element library: Design, features, and insights. *Computers & Mathematics with Applications* **81**, 407–422.
- 915 ARNDT, D., FEHN, N., KANSCHAT, G., KORMANN, K., KRONBICHLER, M., MUNCH, P., WALL, W. A. & WITTE, J. 2020 ExaDG: High-Order Discontinuous Galerkin for the Exa-Scale. In *Software for Exascale Computing - SPPEXA 2016-2019* (ed. H.-J. Bungartz, S. Reiz, B. Uekermann, P. Neumann & W. E. Nagel), pp. 189–224. Cham: Springer International Publishing.
- BEALE, J. T., KATO, T. & MAJDA, A. 1984 Remarks on the breakdown of smooth solutions for the 3-D Euler equations. *Communications in Mathematical Physics* **94** (1), 61–66.
- 920 BORATAV, O. N. & PELZ, R. B. 1994 Direct numerical simulation of transition to turbulence from a high-symmetry initial condition. *Physics of Fluids* **6** (8), 2757–2784.
- BRACHET, M. E. 1991 Direct simulation of three-dimensional turbulence in the Taylor–Green vortex. *Fluid dynamics research* **8** (1-4), 1–8.
- 925 BRACHET, M. E., MEIRON, D. I., ORSZAG, S. A., NICKEL, B. G., MORF, R. H. & FRISCH, U. 1983 Small-scale structure of the Taylor–Green vortex. *Journal of Fluid Mechanics* **130**, 411–452.
- BRACHET, M. E., MENEGUZZI, M., VINCENT, A., POLITANO, H. & SULEM, P. L. 1992 Numerical evidence of smooth self-similar dynamics and possibility of subsequent collapse for three-dimensional ideal flows. *Physics of Fluids A: Fluid Dynamics* **4** (12), 2845–2854.
- 930 BRENIER, Y., DE LELLIS, C. & SZÉKELYHIDI, L. 2011 Weak-strong uniqueness for measure-valued solutions. *Communications in mathematical physics* **305** (2), 351–361.
- BRENNER, M. P., HORMOZ, S. & PUMIR, A. 2016 Potential singularity mechanism for the Euler equations. *Phys. Rev. Fluids* **1**, 084503.
- BROWN, D. L. 1995 Performance of under-resolved two-dimensional incompressible flow simulations. *Journal of Computational Physics* **122** (1), 165 – 183.
- 935 BUCKMASTER, T., DE LELLIS, C. & SZÉKELYHIDI JR., L. 2016 Dissipative Euler flows with Onsager-critical spatial regularity. *Communications on Pure and Applied Mathematics* **69** (9), 1613–1670.
- BUCKMASTER, T., DE LELLIS, C., SZÉKELYHIDI JR., L. & VICOL, V. 2018 Onsager's conjecture for admissible weak solutions. *Communications on Pure and Applied Mathematics* **72** (2), 229–274.
- 940 BUCKMASTER, T., MASMOUDI, N., NOVACK, M. & VICOL, V. 2021 Non-conservative $H^{1/2-}$ weak solutions of the incompressible 3D Euler equations. *arXiv preprint arXiv:2101.09278*.
- BUCKMASTER, T. & VICOL, V. 2019 Nonuniqueness of weak solutions to the Navier–Stokes equation. *Annals of mathematics* **189** (1), 101–144.
- 945 BUCKMASTER, T. & VICOL, V. 2020 Convex integration and phenomenologies in turbulence. *EMS Surveys in Mathematical Sciences* **6** (1), 173–263.
- BURGERS, J.M. 1948 A mathematical model illustrating the theory of turbulence. *Advances in Applied Mechanics*, vol. 1, pp. 171 – 199. Elsevier.
- BUSTAMANTE, M. D. & BRACHET, M. 2012 Interplay between the Beale–Kato–Majda theorem and the analyticity-strip method to investigate numerically the incompressible Euler singularity problem. *Phys. Rev. E* **86**, 066302.
- 950 CAMPOLINA, C. S. & MAILYBAEV, A. A. 2018 Chaotic blowup in the 3D incompressible Euler equations on a logarithmic lattice. *Phys. Rev. Lett.* **121**, 064501.

- CHALMERS, N., AGBAGLAH, G., CHRUST, M. & MAVRIPLIS, C. 2019 A parallel hp-adaptive high order discontinuous Galerkin method for the incompressible Navier–Stokes equations. *Journal of Computational Physics: X* **2**, 100023.
- 955 CHAPELIER, J.-B., DE LA LLAVE PLATA, M. & RENAC, F. 2012 Inviscid and viscous simulations of the Taylor–Green vortex flow using a modal discontinuous Galerkin approach. In *42nd AIAA Fluid Dynamics Conference and Exhibit*, p. 3073.
- CHESKIDOV, A., CONSTANTIN, P., FRIEDLANDER, S. & SHVYDKOY, R. 2008 Energy conservation and Onsager’s conjecture for the Euler equations. *Nonlinearity* **21** (6), 1233–1252.
- 960 CICHOWLAS, C., BONAÏTI, P., DEBBASCH, F. & BRACHET, M. 2005 Effective dissipation and turbulence in spectrally truncated Euler flows. *Phys. Rev. Lett.* **95**, 264502.
- CICHOWLAS, C. & BRACHET, M.-E. 2005 Evolution of complex singularities in Kida–Pelz and Taylor–Green inviscid flows. *Fluid Dynamics Research* **36** (4–6), 239–248.
- 965 CONSTANTIN, P., E, W. & TITI, E. S. 1994 Onsager’s conjecture on the energy conservation for solutions of Euler’s equation. *Communications in Mathematical Physics* **165** (1), 207.
- COPPOLA, G., CAPUANO, F. & DE LUCA, L. 2019 Discrete Energy-Conservation Properties in the Numerical Simulation of the Navier–Stokes Equations. *Applied Mechanics Reviews* **71** (1), 010803.
- DANERI, S., RUNA, E. & SZÉKELYHIDI, L. 2021 Non-uniqueness for the Euler equations up to Onsager’s critical exponent. *Annals of PDE* **7** (1), 1–44.
- 970 DE LELLIS, C. & SZÉKELYHIDI JR., L. 2013 Dissipative continuous Euler flows. *Inventiones mathematicae* **193** (2), 377–407.
- DE LELLIS, C. & SZÉKELYHIDI JR., L. 2014 Dissipative Euler flows and Onsager’s conjecture. *Journal of the European Mathematical Society* **16** (7), 1467–1505.
- 975 DiPERNA, R. J. & MAJDA, A. J. 1987 Oscillations and concentrations in weak solutions of the incompressible fluid equations. *Communications in mathematical physics* **108** (4), 667–689.
- DRIVAS, T. D. & EYINK, G. L. 2019 An onsager singularity theorem for leray solutions of incompressible Navier–Stokes. *Nonlinearity* **32** (11), 4465.
- DRIVAS, T. D. & NGUYEN, H. Q. 2019 Remarks on the emergence of weak Euler solutions in the vanishing viscosity limit. *Journal of Nonlinear Science* **29** (2), 709–721.
- 980 DUBRULLE, B. 2019 Beyond Kolmogorov cascades. *Journal of Fluid Mechanics* **867**, P1.
- DUCHON, J. & ROBERT, R. 2000 Inertial energy dissipation for weak solutions of incompressible Euler and Navier–Stokes equations. *Nonlinearity* **13** (1), 249–255.
- EGGERS, J. 2018 Role of singularities in hydrodynamics. *Phys. Rev. Fluids* **3**, 110503.
- 985 EYINK, G. L. 1994 Energy dissipation without viscosity in ideal hydrodynamics I. Fourier analysis and local energy transfer. *Physica D: Nonlinear Phenomena* **78** (3), 222 – 240.
- EYINK, G. L. 2008 Dissipative anomalies in singular Euler flows. *Physica D: Nonlinear Phenomena* **237** (14), 1956 – 1968, euler Equations: 250 Years On.
- EYINK, G. L. & SREENIVASAN, K. R. 2006 Onsager and the theory of hydrodynamic turbulence. *Rev. Mod. Phys.* **78**, 87–135.
- 990 FEHN, N., HEINZ, J., WALL, W. A. & KRONBICHLER, M. 2021 High-order arbitrary Lagrangian–Eulerian discontinuous Galerkin methods for the incompressible Navier–Stokes equations. *Journal of Computational Physics* **430**, 110040.
- FEHN, N., KRONBICHLER, M., LEHRENFELD, C., LUBE, G. & SCHROEDER, P. W. 2019 High-order DG solvers for under-resolved turbulent incompressible flows: A comparison of L^2 and $H(\text{div})$ methods. *Int. J. Numer. Meth. Fluids* **91** (11), 533–556.
- 995 FEHN, N., MUNCH, P., WALL, W. A. & KRONBICHLER, M. 2020 Hybrid multigrid methods for high-order discontinuous Galerkin discretizations. *Journal of Computational Physics* **415**, 109538.
- FEHN, N., WALL, W. A. & KRONBICHLER, M. 2017 On the stability of projection methods for the incompressible Navier–Stokes equations based on high-order discontinuous Galerkin discretizations. *Journal of Computational Physics* **351**, 392–421.
- 1000 FEHN, N., WALL, W. A. & KRONBICHLER, M. 2018a Efficiency of high-performance discontinuous Galerkin spectral element methods for under-resolved turbulent incompressible flows. *Int. J. Numer. Meth. Fluids* **88** (1), 32–54.
- 1005 FEHN, N., WALL, W. A. & KRONBICHLER, M. 2018b Robust and efficient discontinuous Galerkin methods for under-resolved turbulent incompressible flows. *J. Comput. Phys.* **372**, 667–693.
- FERNANDEZ, P., NGUYEN, N. C. & PERAIRE, J. 2018 On the ability of discontinuous Galerkin methods to simulate under-resolved turbulent flows. *arXiv preprint arXiv:1810.09435* .

- FISCHER, P. & MULLEN, J. 2001 Filter-based stabilization of spectral element methods. *Comptes Rendus de l'Académie des Sciences - Series I - Mathematics* **332** (3), 265 – 270.
- 1010 FLAD, D. & GASSNER, G. 2017 On the use of kinetic energy preserving DG-schemes for large eddy simulation. *Journal of Computational Physics* **350**, 782 – 795.
- FRIGO, M. & JOHNSON, S. G. 2005 The Design and Implementation of FFTW3. *Proceedings of the IEEE* **93** (2), 216–231.
- 1015 FU, G. 2019 An explicit divergence-free DG method for incompressible flow. *Computer Methods in Applied Mechanics and Engineering* **345**, 502 – 517.
- GIBBON, J. D. 2008 The three-dimensional Euler equations: Where do we stand? *Physica D: Nonlinear Phenomena* **237** (14), 1894 – 1904, euler Equations: 250 Years On.
- 1020 GRAFKE, T., HOMANN, H., DREHER, J. & GRAUER, R. 2008 Numerical simulations of possible finite time singularities in the incompressible Euler equations: Comparison of numerical methods. *Physica D: Nonlinear Phenomena* **237** (14), 1932 – 1936, euler Equations: 250 Years On.
- GRAUER, R., MARLIANI, C. & GERMASCHEWSKI, K. 1998 Adaptive mesh refinement for singular solutions of the incompressible Euler equations. *Phys. Rev. Lett.* **80**, 4177–4180.
- 1025 GUZMÁN, J., SHU, C.-W. & SEQUEIRA, F. A. 2016 H(div) conforming and DG methods for incompressible Euler's equations. *IMA Journal of Numerical Analysis* **37** (4), 1733–1771.
- HOU, T. Y. & LI, R. 2008 Blowup or no blowup? The interplay between theory and numerics. *Physica D: Nonlinear Phenomena* **237** (14), 1937 – 1944, euler Equations: 250 Years On.
- 1030 ISETT, P. 2017 Nonuniqueness and existence of continuous, globally dissipative Euler flows. *arXiv preprint arXiv:1710.11186*.
- ISETT, P. 2018 A proof of Onsager's conjecture. *Annals of Mathematics* **188** (3), 871–963.
- JOSSERAND, C., POMEAU, Y. & RICA, S. 2020 Finite-time localized singularities as a mechanism for turbulent dissipation. *Phys. Rev. Fluids* **5**, 054607.
- 1035 KANEDA, Y., ISHIHARA, T., YOKOKAWA, M., ITAKURA, K. & UNO, A. 2003 Energy dissipation rate and energy spectrum in high resolution direct numerical simulations of turbulence in a periodic box. *Physics of Fluids* **15** (2), L21–L24.
- KARNIADAKIS, G. E., ISRAELI, M. & ORSZAG, S. A. 1991 High-order splitting methods for the incompressible Navier–Stokes equations. *J. Comput. Phys.* **97** (2), 414 – 443.
- KARNIADAKIS, G. E. & SHERWIN, S. J. 2005 *Spectral/hp element methods for computational fluid dynamics*, 2nd edn. Oxford University Press.
- 1040 KERR, R. M. 1993 Evidence for a singularity of the three-dimensional, incompressible Euler equations. *Physics of Fluids A: Fluid Dynamics* **5** (7), 1725–1746.
- KERR, R. M. 2013 Bounds for Euler from vorticity moments and line divergence. *Journal of Fluid Mechanics* **729**, R2.
- 1045 KERR, R. M. & HUSSAIN, F. 1989 Simulation of vortex reconnection. *Physica D: Nonlinear Phenomena* **37** (1), 474 – 484.
- KIDA, S. 1985 Three-dimensional periodic flows with high-symmetry. *Journal of the Physical Society of Japan* **54** (6), 2132–2136.
- 1050 KOLMOGOROV, A. N. 1991 The local structure of turbulence in incompressible viscous fluid for very large Reynolds numbers. *Proceedings of the Royal Society of London. Series A: Mathematical and Physical Sciences* **434** (1890), 9–13.
- KRAIS, NICO, SCHNÜCKE, GERO, BOLEMANN, THOMAS & GASSNER, GREGOR J. 2020 Split form ALE discontinuous Galerkin methods with applications to under-resolved turbulent low-Mach number flows. *Journal of Computational Physics* **421**, 109726.
- 1055 KRANK, B., FEHN, N., WALL, W. A. & KRONBICHLER, M. 2017 A high-order semi-explicit discontinuous Galerkin solver for 3D incompressible flow with application to DNS and LES of turbulent channel flow. *J. Comput. Phys.* **348**, 634–659.
- KRONBICHLER, M. & KORMANN, K. 2019 Fast matrix-free evaluation of discontinuous Galerkin finite element operators. *ACM Trans. Math. Softw.* **45** (3), 29:1–29:40.
- 1060 KUZZAY, D., SAW, E.-W., MARTINS, F.J.W.A., FARANDA, D., FOUCAUT, J.-M., DAVIAUD, F. & DUBRULLE, B. 2017 New method for detecting singularities in experimental incompressible flows. *Nonlinearity* **30** (6), 2381.
- LAMBALLAIS, E., DAIRAY, T., LAIZET, S. & VASSILICOS, J. C. 2019 Implicit/explicit spectral viscosity and large-scale SGS effects. In *Direct and Large-Eddy Simulation XI* (ed. Maria Vittoria Salveti, Vincenzo Armenio, Jochen Fröhlich, Bernard J. Geurts & Hans Kuerten), pp. 107–113. Cham: Springer International Publishing.
- 1065

- LARIOS, A., PETERSEN, M. R., TITI, E. S. & WINGATE, B. 2018 A computational investigation of the finite-time blow-up of the 3D incompressible Euler equations based on the Voigt regularization. *Theoretical and Computational Fluid Dynamics* **32** (1), 23–34.
- LIONS, P.-L. 1996 *Mathematical Topics in Fluid Mechanics. Volume 1: Incompressible Models*. Volume 3 of Oxford Lecture Series in Mathematics and its Applications, Oxford University Press.
- 1070 LUO, G. & HOU, T. Y. 2014 Potentially singular solutions of the 3D axisymmetric Euler equations. *Proceedings of the National Academy of Sciences* **111** (36), 12968–12973.
- MANZANERO, J., FERRER, E., RUBIO, G. & VALERO, E. 2020 Design of a Smagorinsky spectral vanishing viscosity turbulence model for discontinuous Galerkin methods. *Computers & Fluids* **200**, 104440.
- 1075 MCKEOWN, RYAN, OSTILLA-MÓNICO, RODOLFO, PUMIR, ALAIN, BRENNER, MICHAEL P. & RUBINSTEIN, SHMUEL M. 2018 Cascade leading to the emergence of small structures in vortex ring collisions. *Phys. Rev. Fluids* **3**, 124702.
- MOFFATT, H. K. 2019 Singularities in fluid mechanics. *Phys. Rev. Fluids* **4**, 110502.
- MORF, R. H., ORSZAG, S. A. & FRISCH, U. 1980 Spontaneous singularity in three-dimensional inviscid, incompressible flow. *Phys. Rev. Lett.* **44**, 572–575.
- 1080 MOURA, R., MENGALDO, G., PEIRÓ, J. & SHERWIN, S. 2017a An LES setting for DG-based implicit LES with insights on dissipation and robustness. In *Spectral and High Order Methods for Partial Differential Equations*. Springer.
- MOURA, R.C., MENGALDO, G., PEIRÓ, J. & SHERWIN, S.J. 2017b On the eddy-resolving capability of high-order discontinuous Galerkin approaches to implicit LES / under-resolved DNS of Euler turbulence. *Journal of Computational Physics* **330**, 615 – 623.
- 1085 MURUGAN, S. D., FRISCH, U., NAZARENKO, S., BESSE, N. & RAY, S. S. 2020 Suppressing thermalization and constructing weak solutions in truncated inviscid equations of hydrodynamics: Lessons from the Burgers equation. *Phys. Rev. Research* **2**, 033202.
- 1090 ONSAGER, L. 1949 Statistical hydrodynamics. *Il Nuovo Cimento (1943-1954)* **6**, 279–287.
- ORLANDI, P. 2009 Energy spectra power laws and structures. *Journal of Fluid Mechanics* **623**, 353–374.
- ORLANDI, P., PIROZZOLI, S. & CARNEVALE, G. F. 2012 Vortex events in Euler and Navier–Stokes simulations with smooth initial conditions. *Journal of Fluid Mechanics* **690**, 288–320.
- PEARSON, B. R., KROGSTAD, P.-Å. & VAN DE WATER, W. 2002 Measurements of the turbulent energy dissipation rate. *Physics of Fluids* **14** (3), 1288–1290.
- 1095 PELZ, R. B. 2001 Symmetry and the hydrodynamic blow-up problem. *Journal of Fluid Mechanics* **444**, 299–320.
- PELZ, R. B. & GULAK, Y. 1997 Evidence for a real-time singularity in hydrodynamics from time series analysis. *Phys. Rev. Lett.* **79**, 4998–5001.
- 1100 PIATKOWSKI, S.-M. 2019 A spectral discontinuous Galerkin method for incompressible flow with applications to turbulence. PhD thesis, Ruprecht-Karls-Universität Heidelberg.
- RAY, S. S., FRISCH, U., NAZARENKO, S. & MATSUMOTO, T. 2011 Resonance phenomenon for the Galerkin-truncated Burgers and Euler equations. *Phys. Rev. E* **84**, 016301.
- SAW, E.-W., KUZZAY, D., FARANDA, D., GUITTONNEAU, A., DAVIAUD, F., WIERTEL-GASQUET, C., PADILLA, V. & DUBRULLE, B. 2016 Experimental characterization of extreme events of inertial dissipation in a turbulent swirling flow. *Nature communications* **7**, 12466.
- 1105 SCHROEDER, P. W. 2019 Robustness of high-order divergence-free finite element methods for incompressible computational fluid dynamics. PhD thesis, Georg-August-Universität Göttingen.
- SCHROEDER, P. W. & LUBE, G. 2018 Divergence-free $H(\text{div})$ -FEM for time-dependent incompressible flows with applications to high Reynolds number vortex dynamics. *Journal of Scientific Computing* **75** (2), 830–858.
- 1110 SHU, C.-W., DON, W.-S., GOTTLIEB, D., SCHILLING, O. & JAMESON, L. 2005 Numerical convergence study of nearly incompressible, inviscid Taylor–Green vortex flow. *Journal of Scientific Computing* **24** (1), 1–27.
- 1115 SREENIVASAN, K. R. 1998 An update on the energy dissipation rate in isotropic turbulence. *Physics of Fluids* **10** (2), 528–529.
- SULEM, C., SULEM, P. L. & FRISCH, H. 1983 Tracing complex singularities with spectral methods. *Journal of Computational Physics* **50** (1), 138 – 161.
- TAYLOR, G. I. 1935 Statistical theory of turbulence. *Proceedings of the Royal Society of London. Series A, Mathematical and Physical Sciences* **151** (873), 421–444.
- 1120 TAYLOR, G. I. & GREEN, A. E. 1937 Mechanism of the production of small eddies from large ones.

Proceedings of the Royal Society of London. Series A, Mathematical and Physical Sciences **158** (895), 499–521.

- ¹¹²⁵ THALABARD, S., BEC, J. & MAILYBAEV, A. A. 2020 From the butterfly effect to spontaneous stochasticity in singular shear flows. *Communications Physics* **3** (1), 1–8.
- WIEDEMANN, EMIL 2017 Weak-strong uniqueness in fluid dynamics. *arXiv preprint arXiv:1705.04220* .
- WINTERS, A. R., MOURA, R. C., MENGALDO, G., GASSNER, G. J., WALCH, S., PEIRO, J. & SHERWIN, S. J. 2018 A comparative study on polynomial dealiasing and split form discontinuous Galerkin schemes for under-resolved turbulence computations. *Journal of Computational Physics* **372**, 1 – 21.